

**HWP1 (Candida) Antibody (internal region)**  
**Peptide-affinity purified goat antibody**  
**Catalog # AF3850a****Specification**

---

**HWP1 (Candida) Antibody (internal region) - Product Information**

|                   |                             |
|-------------------|-----------------------------|
| Application       | E                           |
| Primary Accession | <a href="#">P46593.4</a>    |
| Other Accession   | <a href="#">XP_712998.1</a> |
| Host              | Goat                        |
| Clonality         | Polyclonal                  |
| Concentration     | 0.5 mg/ml                   |
| Isotype           | IgG                         |

**HWP1 (Candida) Antibody (internal region) - Additional Information****Other Names**

HWP1; ECE2

**Format**

0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

HWP1 (Candida) Antibody (internal region) is for research use only and not for use in diagnostic or therapeutic procedures.

**HWP1 (Candida) Antibody (internal region) - Protein Information****HWP1 (Candida) Antibody (internal region) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**HWP1 (Candida) Antibody (internal region) - Images**

**HWP1 (Candida) Antibody (internal region) - References**

Hwp1 and related adhesins contribute to both mating and biofilm formation in *Candida albicans*.  
Ene IV, Bennett RJ. Eukaryot Cell. 2009 Dec;8(12):1909-13. PMID: 19837954