10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

CCAAT/enhancer binding protein zeta (mouse, aa620-633) Antibody (internal region)

Peptide-affinity purified goat antibody
Catalog # AF3888a

Specification

CCAAT/enhancer binding protein zeta (mouse, aa620-633) Antibody (internal region) - Product Information

Application	**WB**
Primary Accession	P53569.2
Other Accession	NP_001019977.1, 12607 (mouse), 362686 (rat)
Reactivity	**Rat**
Predicted	**Mouse, Pig, Dog, Cow**
Host	**Goat**
Clonality	**Polyclonal**
Concentration	**0.5 mg/ml**
Isotype	**IgG**

CCAAT/enhancer binding protein zeta (mouse, aa620-633) Antibody (internal region) - Additional Information

Other Names
Cebpz; CCAAT/enhancer binding protein zeta; AI848081; CBF2; Cbf; Cebpa-rs1; CCAAT-binding factor; CCAAT-box-binding transcription factor; CCAAT/enhancer binding protein alpha (C/EBP) related sequence 1; CCAAT/enhancer-binding protein zeta

Format
0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Storage
Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions
CCAAT/enhancer binding protein zeta (mouse, aa620-633) Antibody (internal region) is for research use only and not for use in diagnostic or therapeutic procedures.

CCAAT/enhancer binding protein zeta (mouse, aa620-633) Antibody (internal region) - Protein Information

CCAAT/enhancer binding protein zeta (mouse, aa620-633) Antibody (internal region) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides

10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

CCAAT/enhancer binding protein zeta (mouse, aa620-633) Antibody (internal region) - Images



AF3888a (1 µg/ml) staining Rat Testis lysate (35 µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

CCAAT/enhancer binding protein zeta (mouse, aa620-633) Antibody (internal region) - References

Upregulation of gene encoding adipogenic transcriptional factors C/EBPalpha and PPARgamma2 in denervated muscle Wagatsuma A. Exp Physiol. 2006 Jul;91(4):747-53 PMID: 16675498