

TBC1D9 Antibody (internal region, near C-Term)
Peptide-affinity purified goat antibody
Catalog # AF4052a**Specification**

TBC1D9 Antibody (internal region, near C-Term) - Product Information

Application	WB
Primary Accession	Q6ZT07
Other Accession	NP_055945.2 , 23158 , 71310 (mouse) , 304645 (rat)
Reactivity	Human
Predicted	Mouse, Rat, Pig, Dog, Cow
Host	Goat
Clonality	Polyclonal
Concentration	0.5 mg/ml
Isotype	IgG
Calculated MW	143229

TBC1D9 Antibody (internal region, near C-Term) - Additional Information**Gene ID** 23158**Other Names**

TBC1 domain family member 9, TBC1 domain family member 9A, TBC1D9, KIAA0882, TBC1D9A

Format

0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

TBC1D9 Antibody (internal region, near C-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

TBC1D9 Antibody (internal region, near C-Term) - Protein Information**Name** TBC1D9**Synonyms** KIAA0882, TBC1D9A**Function**

May act as a GTPase-activating protein for Rab family protein(s).

TBC1D9 Antibody (internal region, near C-Term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

TBC1D9 Antibody (internal region, near C-Term) - Images

AF4052a (0.3 µg/ml) staining of HeLa lysate (35 µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

TBC1D9 Antibody (internal region, near C-Term) - References

Identification and characterization of a novel Tre-2/Bub2/Cdc16 (TBC) protein that possesses Rab3A-GAP activity. Ishibashi K, Kanno E, Itoh T, Fukuda M. Genes Cells. 2009 Jan;14(1):41-52. PMID: 19077034