

syntaxin 11 Antibody (internal region)
Peptide-affinity purified goat antibody
Catalog # AF4065a**Specification**

syntaxin 11 Antibody (internal region) - Product Information

Application	WB, E
Primary Accession	O75558
Other Accession	NP_003755.2 , 8676 , 292483 (rat)
Reactivity	Human
Predicted	Rat, Dog
Host	Goat
Clonality	Polyclonal
Concentration	0.5 mg/ml
Isotype	IgG
Calculated MW	33196

syntaxin 11 Antibody (internal region) - Additional Information**Gene ID** 8676**Other Names**

Syntaxin-11, STX11

Dilution

WB~~1:1000

E~~N/A

Format

0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

syntaxin 11 Antibody (internal region) is for research use only and not for use in diagnostic or therapeutic procedures.

syntaxin 11 Antibody (internal region) - Protein Information**Name** STX11**Function**

SNARE that acts to regulate protein transport between late endosomes and the trans-Golgi network.


Cellular Location

Membrane; Peripheral membrane protein. Golgi apparatus, trans-Golgi network membrane; Peripheral membrane protein

syntaxin 11 Antibody (internal region) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

syntaxin 11 Antibody (internal region) - Images

AF4065a (1 µg/ml) staining of Placenta lysate (35 µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

syntaxin 11 Antibody (internal region) - References

Syntaxin-11, but not syntaxin-2 or syntaxin-4, is required for platelet secretion. Ye S, Karim ZA, Al Hawas R, Pessin JE, Filipovich AH, Whiteheart SW. Blood 2012 Sep 120 (12): 2484-92. PMID: 22767500