

### Goat Anti-HOXA10 Antibody (internal region), Biotinylated

Catalog # AF4299a

### **Specification**

# Goat Anti-HOXA10 Antibody (internal region), Biotinylated - Product Information

Application WB
Primary Accession P31260

Other Accession <u>NP 061824.3</u>, <u>3206</u>

Reactivity Human

Predicted Human, Mouse, Pig, Cow

Host Goat Isotype IgG Calculated MW 42414

## Goat Anti-HOXA10 Antibody (internal region), Biotinylated - Additional Information

#### **Gene ID 3206**

### **Other Names**

Homeobox protein Hox-A10, Homeobox protein Hox-1.8, Homeobox protein Hox-1H, PL, HOXA10, HOX1H

#### Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

## **Precautions**

Goat Anti-HOXA10 Antibody (internal region), Biotinylated is for research use only and not for use in diagnostic or therapeutic procedures.

## Goat Anti-HOXA10 Antibody (internal region), Biotinylated - Protein Information

## Name HOXA10

#### Synonyms HOX1H

### **Function**

Sequence-specific transcription factor which is part of a developmental regulatory system that provides cells with specific positional identities on the anterior-posterior axis. Binds to the DNA sequence 5'-AA[AT]TTTTATTAC-3'.

### **Cellular Location**

Nucleus.

## Goat Anti-HOXA10 Antibody (internal region), Biotinylated - Protocols





Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

# Goat Anti-HOXA10 Antibody (internal region), Biotinylated - Images



Biotinylated AF4299a (0.2 μg/ml) staining of Human Skeletal Muscle lysate (35 μg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.