

Goat Anti-LILRA4 Antibody (C Terminus)
Catalog # AF4307a**Specification****Goat Anti-LILRA4 Antibody (C Terminus) - Product Information**

Application	WB, E
Primary Accession	P59901
Other Accession	NP_036408.3 , 23547
Reactivity	Human
Predicted	Human
Host	Goat
Isotype	IgG
Calculated MW	55181

Goat Anti-LILRA4 Antibody (C Terminus) - Additional Information**Gene ID** 23547**Other Names**

Leukocyte immunoglobulin-like receptor subfamily A member 4, CD85 antigen-like family member G, Immunoglobulin-like transcript 7, ILT-7, CD85g, LILRA4, ILT7

Dilution

WB~~1:1000

E~~N/A

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Goat Anti-LILRA4 Antibody (C Terminus) is for research use only and not for use in diagnostic or therapeutic procedures.

Goat Anti-LILRA4 Antibody (C Terminus) - Protein Information**Name** LILRA4 ([HGNC:15503](#))**Function**

Functions coreceptor to limit the innate immune responses to viral infections; signaling occurs via FCER1G (PubMed:16735691, PubMed:19564354). Down-regulates the production of IFNA1, IFNA2, IFNA4, IFNB1 and TNF by plasmacytoid dendritic cells that have been exposed to influenza virus or cytidine-phosphate-guanosine (CpG) dinucleotides, indicating it functions as a negative regulator of TLR7 and TLR9 signaling cascades (PubMed:16735691, PubMed:16735691, PubMed:16735691).

href="http://www.uniprot.org/citations/19564354" target="_blank">19564354, PubMed:24586760). Down-regulates interferon production in response to interaction with BST2 on HIV-1 infected cells (PubMed:26172439). Activates a signaling cascade in complex with FCER1G that results in phosphorylation of Src family and Syk kinases and thereby triggers mobilization of intracellular Ca(2+) (PubMed:16735691, PubMed:19564354). Does not interfere with the differentiation of plasmacytoid dendritic cells into antigen-presenting cells (PubMed:24586760).

Cellular Location

Cell membrane; Single-pass type I membrane protein

Tissue Location

Detected on plasmacytoid dendritic cells (at protein level). Detected on plasmacytoid dendritic cells, but not on monocytes or B cells.

Goat Anti-LILRA4 Antibody (C Terminus) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Goat Anti-LILRA4 Antibody (C Terminus) - Images



AF4307a (2 µg/ml) staining of Human Spleen lysate (35 µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.