

Goat Anti-CMG1 / CCDC2 / IFT74 Antibody
Peptide-affinity purified goat antibody
Catalog # AF4329a**Specification**

Goat Anti-CMG1 / CCDC2 / IFT74 Antibody - Product Information

Application	IF, FC, E
Primary Accession	O96LB3
Other Accession	NP_001092692.1 , NP_001092693.1 , NP_079379.2 , AAK77221.1
Reactivity	Human
Predicted	Human
Host	Goat
Clonality	Polyclonal
Concentration	100ug/200ul
Isotype	IgG
Calculated MW	69239

Goat Anti-CMG1 / CCDC2 / IFT74 Antibody - Additional Information**Gene ID** 80173**Other Names**

Intraflagellar transport protein 74 homolog, Capillary morphogenesis gene 1 protein, CMG-1, Coiled-coil domain-containing protein 2, IFT74, CCDC2, CMG1

Dilution

E~~1:32000

Format

0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

ImmunogenPeptide with sequence CKTIVDALHSTSGN, from the C or N Terminus of the protein sequence according to [NP_001092692.1](#); [NP_001092693.1](#); [NP_079379.2](#); [AAK77221.1](#).**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Goat Anti-CMG1 / CCDC2 / IFT74 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Goat Anti-CMG1 / CCDC2 / IFT74 Antibody - Protein Information**Name** IFT74

Synonyms CCDC2, CMG1**Function**

Component of the intraflagellar transport (IFT) complex B: together with IFT81, forms a tubulin-binding module that specifically mediates transport of tubulin within the cilium (PubMed:23990561). Binds beta-tubulin via its basic region (PubMed:23990561). Required for ciliogenesis (PubMed:23990561). Essential for flagellogenesis during spermatogenesis (PubMed:33689014).

Cellular Location

Cell projection, cilium. Cytoplasmic vesicle. Cell projection, cilium, flagellum. Cytoplasmic vesicle, secretory vesicle, acrosome {ECO:0000250|UniProtKB:Q8BKE9}. Note=Localizes along primary cilia at interphase and around the basal body/centriole at interphase and mitosis (PubMed:15024030). In male germ cells, strongly expressed in the vesicles of spermatocytes and round spermatids and also in the acrosome and centrosome regions of elongating spermatids and in developing sperm tails (By similarity). {ECO:0000250|UniProtKB:Q8BKE9, ECO:0000269|PubMed:15024030}

Tissue Location


Highly expressed in adult and fetal kidney and expressed at lower level in adult heart, placenta, lung, liver and pancreas, and in fetal heart, lung and liver. Little to no expression was detected in adult brain and skeletal muscle or in fetal brain, thymus and spleen (PubMed:11683410). Detected in sperm (at protein level) (PubMed:33689014).

Goat Anti-CMG1 / CCDC2 / IFT74 Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


Goat Anti-CMG1 / CCDC2 / IFT74 Antibody - Images



AF4329a Immunofluorescence analysis of paraformaldehyde fixed U251 cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (4ug/ml), showing cytoplasmic and Golgi apparatus staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (4ug/ml).



AF4329a Flow cytometric analysis of paraformaldehyde fixed U251 cells (blue line), permeabilized with 0.5% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (0.4ug/ml) showing weak staining. IgG control: Unimmunized goat IgG (black line) followed by Alexa Fluor 488 secondary antibody.



AF4329a (5 µg/ml) staining of paraffin embedded Human Liver. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.