

Goat Anti-PPP1R12A Antibody
Peptide-affinity purified goat antibody
Catalog # AF4330a**Specification**

Goat Anti-PPP1R12A Antibody - Product Information

Application	IF, FC, E
Other Accession	NP_002471.1 , NP_001231919.1 , NP_001231921.1 , NP_001137358.1
Reactivity	Human
Predicted	Human
Host	Goat
Clonality	Polyclonal
Concentration	100ug/200ul
Isotype	IgG

Goat Anti-PPP1R12A Antibody - Additional Information**Dilution**

IF~~1:50~200

FC~~1:10~50

E~~N/A

Format

0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

ImmunogenPeptide with sequence C-REDEYKQKYSRTYD, from the C Terminus of the protein sequence according to [NP_002471.1](#), [NP_001231919.1](#), [NP_001231921.1](#), [NP_001137358.1](#).**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Goat Anti-PPP1R12A Antibody is for research use only and not for use in diagnostic or therapeutic procedures.


Goat Anti-PPP1R12A Antibody - Protein Information**Goat Anti-PPP1R12A Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)

- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


Goat Anti-PPP1R12A Antibody - Images




EB08594 Immunofluorescence analysis of paraformaldehyde fixed A431 cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing cytoplasmic staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).




EB08594 Immunofluorescence analysis of paraformaldehyde fixed U2OS cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing cytoplasmic staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).



EB08594 (4µg/ml) staining of paraffin embedded Human Testis. Heat induced antigen retrieval with citrate buffer pH 6, HRP-staining.



EB08594 Negative Control showing staining of paraffin embedded Human Testis, with no primary antibody.



EB08594 Flow cytometric analysis of paraformaldehyde fixed A431 cells (blue line), permeabilized with 0.5% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (1ug/ml). IgG control: Unimmunized goat IgG (black line) followed by Alexa Fluor 488 secondary antibody.