

Goat anti-AIF1 / IBA1 (isoform 1 and 3), biotinylated Antibody
Peptide-affinity purified goat antibody
Catalog # AF4347a

Specification

Goat anti-AIF1 / IBA1 (isoform 1 and 3), biotinylated Antibody - Product Information

| | |
|-------------------|---|
| Application | WB, Pep-ELISA |
| Primary Accession | P55008 |
| Other Accession | NP_116573.1 , NP_004838.1 , NP_001614.3 |
| Reactivity | Human, Mouse, Rat, Pig |
| Host | Goat |
| Clonality | Polyclonal |
| Calculated MW | 16703 |

Goat anti-AIF1 / IBA1 (isoform 1 and 3), biotinylated Antibody - Additional Information

Gene ID 199

Other Names

AIF1; allograft inflammatory factor 1; AIF-1; IBA1; IRT-1; IRT1; allograft inflammatory factor-1 splice variant Hara-1; interferon gamma responsive transcript; ionized calcium-binding adapter molecule 1; protein G1

Dilution

WB~~1:1000
Pep-ELISA~~N/A

Format

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.

Immunogen

This antibody is expected to recognize isoform 1 (NP_116573.1) and isoform 3 (NP_001614.3).

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Goat anti-AIF1 / IBA1 (isoform 1 and 3), biotinylated Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Goat anti-AIF1 / IBA1 (isoform 1 and 3), biotinylated Antibody - Protein Information

Name AIF1

Synonyms G1, IBA1

Function

Actin-binding protein that enhances membrane ruffling and RAC activation. Enhances the actin-bundling activity of LCP1. Binds calcium. Plays a role in RAC signaling and in phagocytosis. May play a role in macrophage activation and function. Promotes the proliferation of vascular smooth muscle cells and of T-lymphocytes. Enhances lymphocyte migration. Plays a role in vascular inflammation.

Cellular Location

Cytoplasm, cytoskeleton {ECO:0000250|UniProtKB:O70200}. Cell projection, ruffle membrane {ECO:0000250|UniProtKB:O70200}; Peripheral membrane protein {ECO:0000250|UniProtKB:O70200}; Cytoplasmic side {ECO:0000250|UniProtKB:O70200}. Cell projection, phagocytic cup {ECO:0000250|UniProtKB:O70200}. Note=Associated with the actin cytoskeleton at membrane ruffles and at sites of phagocytosis {ECO:0000250|UniProtKB:O70200}

Tissue Location

Detected in T-lymphocytes and peripheral blood mononuclear cells.

Goat anti-AIF1 / IBA1 (isoform 1 and 3), biotinylated Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Goat anti-AIF1 / IBA1 (isoform 1 and 3), biotinylated Antibody - Images