

Goat anti-PARK7 / DJ-1, biotinylated Antibody
Peptide-affinity purified goat antibody
Catalog # AF4356a**Specification**

Goat anti-PARK7 / DJ-1, biotinylated Antibody - Product Information

Application	WB, Pep-ELISA
Primary Accession	O99497
Other Accession	NP_009193.2
Reactivity	Human, Mouse, Rat
Host	Goat
Clonality	Polyclonal
Calculated MW	19891

Goat anti-PARK7 / DJ-1, biotinylated Antibody - Additional Information**Gene ID** 11315**Other Names**

PARK7; parkinson protein 7; DJ-1; DJ1; HEL-S-67p; Parkinson disease (autosomal recessive, early onset) 7; epididymis secretory sperm binding protein Li 67p; oncogene DJ1; protein DJ-1

Dilution

WB~~1:1000

Pep-ELISA~~N/A

Format

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.

Immunogen

Reported variants represent identical protein: NP_009193.2, NP_001116849.1

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Goat anti-PARK7 / DJ-1, biotinylated Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Goat anti-PARK7 / DJ-1, biotinylated Antibody - Protein Information**Name** PARK7 ([HGNC:16369](#))**Function**

Multifunctional protein with controversial molecular function which plays an important role in cell

protection against oxidative stress and cell death acting as oxidative stress sensor and redox-sensitive chaperone and protease (PubMed: [12796482](http://www.uniprot.org/citations/12796482)), PubMed: [17015834](http://www.uniprot.org/citations/17015834)), PubMed: [18711745](http://www.uniprot.org/citations/18711745)), PubMed: [19229105](http://www.uniprot.org/citations/19229105)), PubMed: [20304780](http://www.uniprot.org/citations/20304780)), PubMed: [25416785](http://www.uniprot.org/citations/25416785)), PubMed: [26995087](http://www.uniprot.org/citations/26995087)), PubMed: [28993701](http://www.uniprot.org/citations/28993701)). It is involved in neuroprotective mechanisms like the stabilization of NFE2L2 and PINK1 proteins, male fertility as a positive regulator of androgen signaling pathway as well as cell growth and transformation through, for instance, the modulation of NF-kappa-B signaling pathway (PubMed: [12612053](http://www.uniprot.org/citations/12612053)), PubMed: [14749723](http://www.uniprot.org/citations/14749723)), PubMed: [15502874](http://www.uniprot.org/citations/15502874)), PubMed: [17015834](http://www.uniprot.org/citations/17015834)), PubMed: [18711745](http://www.uniprot.org/citations/18711745)), PubMed: [21097510](http://www.uniprot.org/citations/21097510)). Has been described as a protein and nucleotide deglycase that catalyzes the deglycation of the Maillard adducts formed between amino groups of proteins or nucleotides and reactive carbonyl groups of glyoxals (PubMed: [25416785](http://www.uniprot.org/citations/25416785)), PubMed: [28596309](http://www.uniprot.org/citations/28596309)). But this function is rebutted by other works (PubMed: [27903648](http://www.uniprot.org/citations/27903648)), PubMed: [31653696](http://www.uniprot.org/citations/31653696)). As a protein deglycase, repairs methylglyoxal- and glyoxal-glycated proteins, and releases repaired proteins and lactate or glycolate, respectively. Deglycates cysteine, arginine and lysine residues in proteins, and thus reactivates these proteins by reversing glycation by glyoxals. Acts on early glycation intermediates (hemithioacetals and aminocarbinals), preventing the formation of advanced glycation endproducts (AGE) that cause irreversible damage (PubMed: [25416785](http://www.uniprot.org/citations/25416785)), PubMed: [26995087](http://www.uniprot.org/citations/26995087)), PubMed: [28013050](http://www.uniprot.org/citations/28013050)). Also functions as a nucleotide deglycase able to repair glycated guanine in the free nucleotide pool (GTP, GDP, GMP, dGTP) and in DNA and RNA. Is thus involved in a major nucleotide repair system named guanine glycation repair (GG repair), dedicated to reversing methylglyoxal and glyoxal damage via nucleotide sanitization and direct nucleic acid repair (PubMed: [28596309](http://www.uniprot.org/citations/28596309)). Protects histones from adduction by methylglyoxal, controls the levels of methylglyoxal- derived argininine modifications on chromatin (PubMed: [30150385](http://www.uniprot.org/citations/30150385)). Able to remove the glycations and restore histone 3, histone glycation disrupts both local and global chromatin architecture by altering histone-DNA interactions as well as histone acetylation and ubiquitination levels (PubMed: [30150385](http://www.uniprot.org/citations/30150385)), PubMed: [30894531](http://www.uniprot.org/citations/30894531)). Displays a very low glyoxalase activity that may reflect its deglycase activity (PubMed: [22523093](http://www.uniprot.org/citations/22523093)), PubMed: [28993701](http://www.uniprot.org/citations/28993701)), PubMed: [31653696](http://www.uniprot.org/citations/31653696)). Eliminates hydrogen peroxide and protects cells against hydrogen peroxide-induced cell death (PubMed: [16390825](http://www.uniprot.org/citations/16390825)). Required for correct mitochondrial morphology and function as well as for autophagy of dysfunctional mitochondria (PubMed: [16632486](http://www.uniprot.org/citations/16632486)), PubMed: [19229105](http://www.uniprot.org/citations/19229105)). Plays a role in regulating expression or stability of the mitochondrial uncoupling proteins SLC25A14 and SLC25A27 in dopaminergic neurons of the

substantia nigra pars compacta and attenuates the oxidative stress induced by calcium entry into the neurons via L-type channels during pacemaking (PubMed:18711745). Regulates astrocyte inflammatory responses, may modulate lipid rafts-dependent endocytosis in astrocytes and neuronal cells (PubMed:23847046). In pancreatic islets, involved in the maintenance of mitochondrial reactive oxygen species (ROS) levels and glucose homeostasis in an age- and diet dependent manner. Protects pancreatic beta cells from cell death induced by inflammatory and cytotoxic setting (By similarity). Binds to a number of mRNAs containing multiple copies of GG or CC motifs and partially inhibits their translation but dissociates following oxidative stress (PubMed:18626009). Metal-binding protein able to bind copper as well as toxic mercury ions, enhances the cell protection mechanism against induced metal toxicity (PubMed:23792957). In macrophages, interacts with the NADPH oxidase subunit NCF1 to direct NADPH oxidase-dependent ROS production, and protects against sepsis (By similarity).

Cellular Location

Cell membrane {ECO:0000250|UniProtKB:Q99LX0}; Lipid-anchor {ECO:0000250|UniProtKB:Q99LX0}. Cytoplasm. Nucleus. Membrane raft {ECO:0000250|UniProtKB:O88767}. Mitochondrion. Endoplasmic reticulum. Note=Under normal conditions, located predominantly in the cytoplasm and, to a lesser extent, in the nucleus and mitochondrion. Translocates to the mitochondrion and subsequently to the nucleus in response to oxidative stress and exerts an increased cytoprotective effect against oxidative damage (PubMed:18711745). Detected in tau inclusions in brains from neurodegenerative disease patients (PubMed:14705119). Membrane raft localization in astrocytes and neuronal cells requires palmitoylation

Tissue Location

Highly expressed in pancreas, kidney, skeletal muscle, liver, testis and heart. Detected at slightly lower levels in placenta and brain (at protein level). Detected in astrocytes, Sertoli cells, spermatogonia, spermatids and spermatozoa. Expressed by pancreatic islets at higher levels than surrounding exocrine tissues (PubMed:22611253).

Goat anti-PARK7 / DJ-1, biotinylated Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Goat anti-PARK7 / DJ-1, biotinylated Antibody - Images