

Goat anti-FHL1 / SLIM1, Biotinylated Antibody
Peptide-affinity purified goat antibody
Catalog # AF4384a

Specification

Goat anti-FHL1 / SLIM1, Biotinylated Antibody - Product Information

Application	WB, IHC, Pep-ELISA
Primary Accession	Q13642
Other Accession	NP_001440.2 , NP_001153171.1
Reactivity	Human, Mouse, Rat, Pig, Dog, Bovine
Host	Goat
Clonality	Polyclonal
Calculated MW	36263

Goat anti-FHL1 / SLIM1, Biotinylated Antibody - Additional Information

Gene ID 2273

Other Names

FHL1; four and a half LIM domains 1; FHL-1; FHL1A; FHL1B; FLH1A; KYOT; SLIM; SLIM-1; SLIM1; SLIMMER; XMPMA; LIM protein SLIMMER; four-and-a-half Lin11, Isl-1 and Mec-3 domains 1; skeletal muscle LIM-protein 1

Dilution

WB~~1:1000
IHC~~1:100~500
Pep-ELISA~~N/A

Format

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.

Immunogen

This antibody is expected to recognise isoforms 2 (NP_001440.1) and 5 (NP_001153171.1). Since June 2009 the second N residue in the immunizing peptide is no longer representative as it has been substituted by Q at position 284 of NP_001440.2. Variants NP

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Goat anti-FHL1 / SLIM1, Biotinylated Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Goat anti-FHL1 / SLIM1, Biotinylated Antibody - Protein Information

Name FHL1

Synonyms SLIM1

Function

May have an involvement in muscle development or hypertrophy.

Cellular Location

[Isoform 1]: Cytoplasm. [Isoform 2]: Nucleus. Cytoplasm, cytosol. Note=Predominantly nuclear in myoblasts but is cytosolic in differentiated myotubes

Tissue Location

Isoform 1 is highly expressed in skeletal muscle and to a lesser extent in heart, placenta, ovary, prostate, testis, small intestine, colon and spleen. Expression is barely detectable in brain, lung, liver, kidney, pancreas, thymus and peripheral blood leukocytes. Isoform 2 is expressed in brain, skeletal muscle and to a lesser extent in heart, colon, prostate and small intestine. Isoform 3 is expressed in testis, heart and skeletal muscle

Goat anti-FHL1 / SLIM1, Biotinylated Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Goat anti-FHL1 / SLIM1, Biotinylated Antibody - Images