

Goat anti-SSP29 / ANP32B, Biotinylated Antibody

Peptide-affinity purified goat antibody Catalog # AF4385a

Specification

Goat anti-SSP29 / ANP32B, Biotinylated Antibody - Product Information

Application Primary Accession Other Accession Reactivity Host

Clonality Calculated MW WB, IHC, Pep-ELISA

Q92688 NP_006392.1

Human, Mouse, Rat, Pig, Dog, Bovine

Goat Polyclonal 28788

Goat anti-SSP29 / ANP32B, Biotinylated Antibody - Additional Information

Gene ID 10541

Other Names

ANP32B; acidic nuclear phosphoprotein 32 family member B; APRIL; PHAPI2; SSP29; acidic (leucine-rich) nuclear phosphoprotein 32 family, member B; acidic protein rich in leucines; putative HLA-DR-associated protein I-2; silver-stainable protein SSP29

Dilution

WB~~1:1000 IHC~~1:100~500 Pep-ELISA~~N/A

Format

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.

Immunogen

This antibody does NOT recognize the TNF family member also known as APRIL - Gene ID number 8741. However this antibody may cross-react with ANP32A (GeneID 8125).

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Goat anti-SSP29 / ANP32B, Biotinylated Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Goat anti-SSP29 / ANP32B, Biotinylated Antibody - Protein Information

Name ANP32B



Synonyms APRIL, PHAPI2

Function

Multifunctional protein that is involved in the regulation of many processes including cell proliferation, apoptosis, cell cycle progression or transcription (PubMed: 18039846, PubMed:20015864). Regulates the proliferation of neuronal stem cells, differentiation of leukemic cells and progression from G1 to S phase of the cell cycle. As negative regulator of caspase-3-dependent apoptosis, may act as an antagonist of ANP32A in regulating tissue homeostasis (PubMed: 20015864). Exhibits histone chaperone properties, able to recruit histones to certain promoters, thus regulating the transcription of specific genes (PubMed: 18039846, PubMed:20538007). Also plays an essential role in the nucleocytoplasmic transport of specific mRNAs via the uncommon nuclear mRNA export receptor XPO1/CRM1 (PubMed: 17178712). Participates in the regulation of adequate adaptive immune responses by acting on mRNA expression and cell proliferation (By similarity).

Cellular Location

[Isoform 1]: Nucleus. Cytoplasm Note=Accumulates in the nuclei at the S phase.

Tissue Location

Expressed in heart, lung, pancreas, prostate and in spleen, thymus and placenta.

Goat anti-SSP29 / ANP32B, Biotinylated Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Goat anti-SSP29 / ANP32B, Biotinylated Antibody - Images