

Goat anti-58K Golgi protein(N-Term)/FTCD, Biotinylated Antibody
Peptide-affinity purified goat antibody
Catalog # AF4387a**Specification**

Goat anti-58K Golgi protein(N-Term)/FTCD, Biotinylated Antibody - Product Information

Application	WB, IHC, Pep-ELISA
Primary Accession	O95954
Other Accession	NP_006648.1
Reactivity	Human, Mouse, Pig
Host	Goat
Clonality	Polyclonal
Calculated MW	58927

Goat anti-58K Golgi protein(N-Term)/FTCD, Biotinylated Antibody - Additional Information**Gene ID** 10841**Other Names**

FTCD; formimidoyltransferase cyclodeaminase; LCHC1; formiminotransferase cyclodeaminase; formiminotransferase-cyclodeaminase

DilutionWB~~1:1000
IHC~~1:100~500
Pep-ELISA~~N/A**Format**

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.

Immunogen

Reported variants represent identical protein: NP_996848.1, NP_006648.1 Reported variants represent identical protein: XP_006724024.2, XP_011527739.1

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Goat anti-58K Golgi protein(N-Term)/FTCD, Biotinylated Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Goat anti-58K Golgi protein(N-Term)/FTCD, Biotinylated Antibody - Protein Information**Name** FTCD

Function

Folate-dependent enzyme, that displays both transferase and deaminase activity. Serves to channel one-carbon units from formiminoglutamate to the folate pool.

Cellular Location

Cytoplasm, cytosol {ECO:0000250|UniProtKB:Q9YH58}. Golgi apparatus {ECO:0000250|UniProtKB:Q9YH58}. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome, centriole Note=More abundantly located around the mother centriole

Goat anti-58K Golgi protein(N-Term)/FTCD, Biotinylated Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Goat anti-58K Golgi protein(N-Term)/FTCD, Biotinylated Antibody - Images