

Goat anti-Latexin, Biotinylated Antibody
Peptide-affinity purified goat antibody
Catalog # AF4389a**Specification**

Goat anti-Latexin, Biotinylated Antibody - Product Information

| | |
|-------------------|-----------------------------|
| Application | WB, IHC, IF, Pep-ELISA |
| Primary Accession | Q9BS40 |
| Other Accession | NP_064554.3 |
| Reactivity | Human, Mouse, Rat, Dog |
| Host | Goat |
| Clonality | Polyclonal |
| Calculated MW | 25750 |

Goat anti-Latexin, Biotinylated Antibody - Additional Information**Gene ID** 56925**Other Names**

LXN; latexin; ECI; TCI; MUM; endogenous carboxypeptidase inhibitor; tissue carboxypeptidase inhibitor

DilutionWB~~1:1000
IHC~~1:100~500
IF~~1:50~200
Pep-ELISA~~N/A**Format**

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Goat anti-Latexin, Biotinylated Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Goat anti-Latexin, Biotinylated Antibody - Protein Information**Name** LXN**Function**

Hardly reversible, non-competitive, and potent inhibitor of CPA1, CPA2 and CPA4. May play a role in inflammation.

Cellular Location

Cytoplasm.

Tissue Location

Highly expressed in heart, prostate, ovary, kidney, pancreas, and colon, moderate or low in other tissues including brain

Goat anti-Latexin, Biotinylated Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Goat anti-Latexin, Biotinylated Antibody - Images