

Goat anti-CSK, Biotinylated Antibody
Peptide-affinity purified goat antibody
Catalog # AF4435a**Specification**

Goat anti-CSK, Biotinylated Antibody - Product Information

Application	WB, IHC, Pep-ELISA
Primary Accession	P41240
Other Accession	NP_004374.1
Reactivity	Human, Mouse, Rat, Dog, Bovine
Host	Goat
Clonality	Polyclonal
Calculated MW	50704

Goat anti-CSK, Biotinylated Antibody - Additional Information**Gene ID** 1445**Other Names**

CSK; c-src tyrosine kinase; C-Src kinase; protein-tyrosine kinase CYL

Format

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.

Immunogen

Reported variants represent identical protein: NP_004374.1, NP_001120662.1

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Goat anti-CSK, Biotinylated Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Goat anti-CSK, Biotinylated Antibody - Protein Information**Name** CSK**Function**

Non-receptor tyrosine-protein kinase that plays an important role in the regulation of cell growth, differentiation, migration and immune response. Phosphorylates tyrosine residues located in the C-terminal tails of Src-family kinases (SFKs) including LCK, SRC, HCK, FYN, LYN, CSK or YES1. Upon tail phosphorylation, Src-family members engage in intramolecular interactions between the phosphotyrosine tail and the SH2 domain that result in an inactive conformation. To inhibit SFKs, CSK is recruited to the plasma membrane via binding to transmembrane proteins or adapter

proteins located near the plasma membrane. Suppresses signaling by various surface receptors, including T-cell receptor (TCR) and B-cell receptor (BCR) by phosphorylating and maintaining inactive several positive effectors such as FYN or LCK.

Cellular Location

Cytoplasm. Cell membrane. Note=Mainly cytoplasmic, also present in lipid rafts

Tissue Location

Expressed in lung and macrophages.

Goat anti-CSK, Biotinylated Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Goat anti-CSK, Biotinylated Antibody - Images