

## KD-Validated Anti-ITPK1 Rabbit Monoclonal Antibody

### Rabbit monoclonal antibody

Catalog # AGI1042

### Specification

#### KD-Validated Anti-ITPK1 Rabbit Monoclonal Antibody - Product Information

Application	WB, FC, ICC
Primary Accession	<a href="#">Q13572</a>
Reactivity	Human
Clonality	Monoclonal
Isotype	Rabbit IgG
Calculated MW	Predicted, 46 kDa, observed, 42 kDa kDa
Gene Name	ITPK1
Aliases	ITPK1; Inositol-Tetrakisphosphate 1-Kinase; Inositol 1,3,4-Trisphosphate 5/6-Kinase; Ins(1,3,4)P(3) 5/6-Kinase; Inositol 1,3,4-Triphosphate 5/6 Kinase; Inositol-Triphosphate 5/6-Kinase; EC 2.7.1.134; EC 2.7.1.159; ITRPK1
Immunogen	A synthesized peptide derived from human ITPK1

#### KD-Validated Anti-ITPK1 Rabbit Monoclonal Antibody - Additional Information

Gene ID	3705
Other Names	Inositol-tetrakisphosphate 1-kinase, 2.7.1.134, Inositol 1, 3, 4-trisphosphate 5/6-kinase, Inositol-triphosphate 5/6-kinase, Ins(1, 3, 4)P(3) 5/6-kinase, 2.7.1.159, ITPK1 ( <a href="http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=6177" target="_blank">http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=6177</a> )

#### KD-Validated Anti-ITPK1 Rabbit Monoclonal Antibody - Protein Information

Name ITPK1 ([HGNC:6177](#))

#### Function

Kinase that can phosphorylate various inositol polyphosphate such as Ins(3,4,5,6)P4 or Ins(1,3,4)P3 (PubMed: [11042108](http://www.uniprot.org/citations/11042108), PubMed: [8662638](http://www.uniprot.org/citations/8662638)). Phosphorylates Ins(3,4,5,6)P4 at position 1 to form Ins(1,3,4,5,6)P5 (PubMed: [11042108](http://www.uniprot.org/citations/11042108)). This reaction is thought to have regulatory importance, since Ins(3,4,5,6)P4 is an inhibitor of plasma membrane Ca(2+)-activated Cl(-) channels, while Ins(1,3,4,5,6)P5 is not. Also phosphorylates Ins(1,3,4)P3 on O-5 and O-6 to form Ins(1,3,4,6)P4, an essential molecule in the hexakisphosphate (InsP6) pathway (PubMed: [11042108](http://www.uniprot.org/citations/11042108), PubMed: [8662638](http://www.uniprot.org/citations/8662638)). Also acts as an inositol polyphosphate phosphatase that dephosphorylates Ins(1,3,4,5)P4 and Ins(1,3,4,6)P4 to

Ins(1,3,4)P<sub>3</sub>, and Ins(1,3,4,5,6)P<sub>5</sub> to Ins(3,4,5,6)P<sub>4</sub> (PubMed:<a href="http://www.uniprot.org/citations/11909533" target="\_blank">11909533</a>, PubMed:<a href="http://www.uniprot.org/citations/17616525" target="\_blank">17616525</a>). May also act as an isomerase that interconverts the inositol tetrakisphosphate isomers Ins(1,3,4,5)P<sub>4</sub> and Ins(1,3,4,6)P<sub>4</sub> in the presence of ADP and magnesium (PubMed:<a href="http://www.uniprot.org/citations/11909533" target="\_blank">11909533</a>). Probably acts as the rate-limiting enzyme of the InsP<sub>6</sub> pathway. Modifies TNF- $\alpha$ -induced apoptosis by interfering with the activation of TNFRSF1A-associated death domain (PubMed:<a href="http://www.uniprot.org/citations/11909533" target="\_blank">11909533</a>, PubMed:<a href="http://www.uniprot.org/citations/12925536" target="\_blank">12925536</a>, PubMed:<a href="http://www.uniprot.org/citations/17616525" target="\_blank">17616525</a>). Plays an important role in MLKL-mediated necroptosis. Produces highly phosphorylated inositol phosphates such as inositolhexakisphosphate (InsP<sub>6</sub>) which bind to MLKL mediating the release of an N-terminal auto-inhibitory region leading to its activation. Essential for activated phospho-MLKL to oligomerize and localize to the cell membrane during necroptosis (PubMed:<a href="http://www.uniprot.org/citations/17616525" target="\_blank">17616525</a>).

### Tissue Location


Expressed in brain > heart > skeletal muscle = kidney = pancreas = liver = placenta > lung. In brain, it is expressed in cerebellum, cerebral cortex, medulla, spinal cord, occipital lobe, frontal lobe, temporal lobe and putamen.

### KD-Validated Anti-ITPK1 Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


### KD-Validated Anti-ITPK1 Rabbit Monoclonal Antibody - Images



Copyright ©2025 Genuin Biotechnologies LLC


Western blotting analysis using anti-ITPK1 antibody (Cat#AGI1042). Total cell lysates (30  $\mu$ g) from various cell lines were loaded and separated by SDS-PAGE. The blot was incubated with

anti-ITPK1 antibody (Cat#AGI1042, 1:5,000) and HRP-conjugated goat anti-rabbit secondary antibody respectively.




Copyright ©2025 Genuin Biotechnologies LLC

Western blotting analysis using anti-ITPK1 antibody (Cat#AGI1042). ITPK1 expression in wild type (WT) and ITPK1 shRNA knockdown (KD) HeLa cells with 20 µg of total cell lysates. Hsp90 α serves as a loading control. The blot was incubated with anti-ITPK1 antibody (Cat#AGI1042, 1:5,000) and HRP-conjugated goat anti-rabbit secondary antibody respectively.



Copyright ©2024 Genuin Biotechnologies LLC

Flow cytometric analysis of ITPK1 expression in HepG2 cells using ITPK1 antibody (Cat#AGI1042, 1:2,000). Green, isotype control; red, ITPK1.



Immunocytochemical staining of HepG2 cells with anti-ITPK1 antibody (Cat#AGI1042, 1:1,000). Nuclei were stained blue with DAPI; ITPK1 was stained magenta with Alexa Fluor® 647. Images were taken using Leica stellaris 5. Protein abundance based on laser Intensity and smart gain: Medium. Scale bar: 20 µm.