

#### KD-Validated Anti-TXNDC5 Mouse Monoclonal Antibody Mouse monoclonal antibody Catalog # AGI1114

### **Specification**

# **KD-Validated Anti-TXNDC5 Mouse Monoclonal Antibody - Product Information**

Aliases TXNDC5; Thioredoxin Domain Containing ERp46; EndoPDI; PDIA15; Hcc-2; Protein Disulfide Isomerase Family A, Member 15 Endoplasmic Reticulum Resident Protein 46; Thioredoxin Domain-Containing Prote 5; Thioredoxin-Like Protein P46; ER Prote 46; FLJ21353; FLJ90810; MGC3178; Thioredoxin Domain Containing 5 (Endoplasmic Reticulum); Epididymis Secretory Sperm Binding Protein; Endothelial Protein Disulphide Isomerase Endoplasmic Reticulum Protein ERp46; Thioredoxin Related Protein; EC 1.8.4; E 5.3.4.1; UNQ364; STRF8; TLP46; HCC2	5; ein ein
Immunogen Recombinant protein of human TXNDC5	

### KD-Validated Anti-TXNDC5 Mouse Monoclonal Antibody - Additional Information

Gene ID 81567 Other Names Thioredoxin domain-containing protein 5, Thioredoxin-like protein p46, TXNDC5 (<a href="http://www.genenames.org/cgi-bin/gene\_symbol\_report?hgnc\_id=21073" target="\_blank">HGNC:21073</a>), TLP46

### **KD-Validated Anti-TXNDC5 Mouse Monoclonal Antibody - Protein Information**

Name TXNDC5 (HGNC:21073)

Synonyms TLP46

Function

Protein disulfide isomerase of the endoplasmic reticulum lumen involved in the formation of disulfide bonds in proteins. Can reduce insulin disulfide bonds.

**Cellular Location** 




Endoplasmic reticulum lumen {ECO:0000250|UniProtKB:Q91W90, ECO:0000255|PROSITE-ProRule:PRU10138}

# **KD-Validated Anti-TXNDC5 Mouse Monoclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>


**KD-Validated Anti-TXNDC5 Mouse Monoclonal Antibody - Images** 



Western blotting analysis using anti-thioredoxin domain containing 5 antibody (Cat#AGI1114). Total cell lysates ( $30 \mu g$ ) from various cell lines were loaded and separated by SDS-PAGE. The blot was incubated with anti-thioredoxin domain containing 5 antibody (Cat#AGI1114, 1:5,000) and HRP-conjugated goat anti-mouse secondary antibody respectively.

kDa	N 10	kDa	n 40
250 -		250 -	
150 -		150 —	
100 <b>—</b> 75 <b>—</b>		Hsp90 α 75 -	
75-			
50 <b>—</b>		50 <b>—</b>	Thioredoxin domain containing 5
37 -		37 -	
0.			
25 <b>—</b>		25 -	
20 —		20 -	

Western blotting analysis using anti-thioredoxin domain containing 5 antibody (Cat#AGI1114). Thioredoxin domain containing 5 expression in wild type (WT) and thioredoxin domain containing 5 (TXNDC5) knockdown (KD) HSHC cells with 30  $\mu$ g of total cell lysates. Hsp90  $\alpha$  serves as a loading control. The blot was incubated with anti-thioredoxin domain containing 5 antibody (Cat#AGI1114, 1:2,500) and HRP-conjugated goat anti-mouse secondary antibody respectively.



Thioredoxin domain containing 5-Alexa Fluor® 647

Flow cytometric analysis of Thioredoxin domain containing 5 expression in HepG2 cells using anti-Thioredoxin domain containing 5 antibody (Cat# AGI1114, 1:2,000). Green, isotype control; red, Thioredoxin domain containing 5.