

KD-Validated Anti-AP2S1 Rabbit Monoclonal Antibody
Rabbit monoclonal antibody
Catalog # AGI1149**Specification****KD-Validated Anti-AP2S1 Rabbit Monoclonal Antibody - Product Information**

Application	WB, FC, ICC
Primary Accession	P53680
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Isotype	Rabbit IgG
Calculated MW	Predicted, 17 kDa , observed, 17 kDa
Gene Name	KDa AP2S1
Aliases	AP2S1; Adaptor Related Protein Complex 2 Subunit Sigma 1; CLAPS2; FBH3; Adaptor Related Protein Complex 2 Sigma 1 Subunit; Clathrin Assembly Protein 2 Sigma Small Chain; Plasma Membrane Adaptor AP-2 17 KDa Protein; Adaptor Protein Complex AP-2 Subunit Sigma; Clathrin Coat-Associated Protein AP17; Clathrin Coat Assembly Protein AP17; AP-2 Complex Subunit Sigma; HA2 17 KDa Subunit; Sigma2-Adaptin; Sigma-2; FBHOk; AP17; HHC3; Clathrin-Associated/Assembly/Adaptor Protein, Small 2 (17kD); Adaptor-Related Protein Complex 2 Subunit Sigma; Hypocalciuric Hypercalcemia 3 (Oklahoma Type); FBHOK
Immunogen	A synthesized peptide derived from AP2S1

KD-Validated Anti-AP2S1 Rabbit Monoclonal Antibody - Additional Information

Gene ID	1175
Other Names	AP-2 complex subunit sigma, Adaptor protein complex AP-2 subunit sigma, Adaptor-related protein complex 2 subunit sigma, Clathrin assembly protein 2 sigma small chain, Clathrin coat assembly protein AP17, Clathrin coat-associated protein AP17, HA2 17 kDa subunit, Plasma membrane adaptor AP-2 17 kDa protein, Sigma2-adaptin, AP2S1 (http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=565)
	target="_blank">HGNC:565), AP17, CLAPS2

KD-Validated Anti-AP2S1 Rabbit Monoclonal Antibody - Protein Information**Name** AP2S1 ([HGNC:565](#))**Synonyms** AP17, CLAPS2

Function

Component of the adaptor protein complex 2 (AP-2). Adaptor protein complexes function in protein transport via transport vesicles in different membrane traffic pathways. Adaptor protein complexes are vesicle coat components and appear to be involved in cargo selection and vesicle formation. AP-2 is involved in clathrin-dependent endocytosis in which cargo proteins are incorporated into vesicles surrounded by clathrin (clathrin-coated vesicles, CCVs) which are destined for fusion with the early endosome. The clathrin lattice serves as a mechanical scaffold but is itself unable to bind directly to membrane components. Clathrin-associated adaptor protein (AP) complexes which can bind directly to both the clathrin lattice and to the lipid and protein components of membranes are considered to be the major clathrin adaptors contributing the CCV formation. AP-2 also serves as a cargo receptor to selectively sort the membrane proteins involved in receptor-mediated endocytosis. AP-2 seems to play a role in the recycling of synaptic vesicle membranes from the presynaptic surface. AP-2 recognizes Y-X-X-[FILMV] (Y-X-X-Phi) and [ED]-X-X-X-L- [LI] endocytosis signal motifs within the cytosolic tails of transmembrane cargo molecules. AP-2 may also play a role in maintaining normal post-endocytic trafficking through the ARF6-regulated, non- clathrin pathway. The AP-2 alpha and AP-2 sigma subunits are thought to contribute to the recognition of the [ED]-X-X-X-L-[LI] motif (By similarity). May also play a role in extracellular calcium homeostasis.


Cellular Location


Cell membrane {ECO:0000250|UniProtKB:P63010}. Membrane, coated pit; Peripheral membrane protein; Cytoplasmic side. Note=AP-2 appears to be excluded from internalizing CCVs and to disengage from sites of endocytosis seconds before internalization of the nascent CCV {ECO:0000250|UniProtKB:P63010}

KD-Validated Anti-AP2S1 Rabbit Monoclonal Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


KD-Validated Anti-AP2S1 Rabbit Monoclonal Antibody - Images




Western blotting analysis using anti-AP2S1 antibody (Cat#AGI1149). Total cell lysates (30 μ g) from various cell lines were loaded and separated by SDS-PAGE. The blot was incubated with anti-AP2S1 antibody (Cat#AGI1149, 1:5,000) and HRP-conjugated goat anti-rabbit secondary antibody respectively.



Western blotting analysis using anti-AP2S1 antibody (Cat#AGI1149). AP2S1 expression in wild type (WT) and AP2S1 shRNA knockdown (KD) HeLa cells with 30 μ g of total cell lysates. β -Tubulin serves as a loading control. The blot was incubated with anti-AP2S1 antibody (Cat#AGI1149 1:5,000) and HRP-conjugated goat anti-rabbit secondary antibody respectively.



Flow cytometric analysis of AP2S1 expression in C2C12 cells using AP2S1 antibody (Cat#AGI1149, 1:2,000). Green, isotype control; red, AP2S1.



Immunocytochemical staining of C2C12 cells with AP2S1 antibody (Cat#AGI1149, 1:1,000). Nuclei were stained blue with DAPI; AP2S1 was stained magenta with Alexa Fluor® 647. Images were taken using Leica stellaris 5. Protein abundance based on laser Intensity and smart gain: Medium. Scale bar: 20 μ m.