

KD-Validated Anti-Androgen receptor Rabbit Monoclonal Antibody
Rabbit monoclonal antibody
Catalog # AGI1162**Specification****KD-Validated Anti-Androgen receptor Rabbit Monoclonal Antibody - Product Information**

Application	WB, ICC
Primary Accession	P10275
Reactivity	Human, Mouse
Clonality	Monoclonal
Isotype	Rabbit IgG
Calculated MW	Predicted, 99 kDa , observed, 110 kDa
Gene Name	KDa
Aliases	AR AR; Androgen Receptor; NR3C4; Dihydrotestosterone Receptor; HUMARA; SMA1; DHTR; AIS; Nuclear Receptor Subfamily 3 Group C Member; SBMA; Spinal And Bulbar Muscular Atrophy; Testicular; Feminization; Kennedy Disease; HSP1; AR8; TFM; KD
Immunogen	A synthesized peptide derived from human Androgen Receptor

KD-Validated Anti-Androgen receptor Rabbit Monoclonal Antibody - Additional Information

Gene ID	367
Other Names	
Androgen receptor, Dihydrotestosterone receptor, Nuclear receptor subfamily 3 group C member 4, AR, DHTR, NR3C4	

KD-Validated Anti-Androgen receptor Rabbit Monoclonal Antibody - Protein Information**Name** AR**Synonyms** DHTR, NR3C4**Function**

Steroid hormone receptors are ligand-activated transcription factors that regulate eukaryotic gene expression and affect cellular proliferation and differentiation in target tissues (PubMed:19022849). Transcription factor activity is modulated by bound coactivator and corepressor proteins like ZBTB7A that recruits NCOR1 and NCOR2 to the androgen response elements/ARE on target genes, negatively regulating androgen receptor signaling and androgen-induced cell proliferation (PubMed:20812024). Transcription activation is also down-regulated by NR0B2. Activated, but not phosphorylated, by HIPK3 and ZIPK/DAPK3.

Cellular Location

Nucleus. Cytoplasm Note=Detected at the promoter of target genes (PubMed:25091737)
Predominantly cytoplasmic in unligated form but translocates to the nucleus upon ligand-binding.
Can also translocate to the nucleus in unligated form in the presence of RACK1.

Tissue Location


[Isoform 2]: Mainly expressed in heart and skeletal muscle.

KD-Validated Anti-Androgen receptor Rabbit Monoclonal Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


KD-Validated Anti-Androgen receptor Rabbit Monoclonal Antibody - Images



Western blotting analysis using anti-Androgen receptor antibody (Cat#AGI1162). Total cell lysates (30 µg) from various cell lines were loaded and separated by SDS-PAGE. The blot was incubated with anti-Androgen receptor antibody (Cat#AGI1162, 1:5,000) and HRP-conjugated goat anti-rabbit secondary antibody respectively.



Western blotting analysis using anti-Androgen receptor antibody (Cat#AGI1162). Androgen receptor expression in wild type (WT) and Androgen receptor shRNA knockdown (KD) HeLa cells with 30 µg of total cell lysates. β-Tubulin serves as a loading control. The blot was incubated with anti-Androgen receptor antibody (Cat#AGI1162, 1:5,000) and HRP-conjugated goat anti-rabbit secondary antibody respectively.



Immunocytochemical staining of C2C12 cells with Androgen receptor antibody (Cat#AG1162, 1:1,000). Nuclei were stained blue with DAPI; Androgen receptor was stained magenta with Alexa Fluor® 647. Images were taken using Leica stellaris 5. Protein abundance based on laser Intensity and smart gain: Low. Scale bar: 20 μ m.