

KD-Validated Anti-DDX5 Rabbit Monoclonal Antibody Rabbit monoclonal antibody Catalog # AGI1197

Specification

KD-Validated Anti-DDX5 Rabbit Monoclonal Antibody - Product Information

WB, FC, ICC
<u>P17844</u>
Rat, Human, Mouse
Monoclonal
Rabbit IgG
Predicted, 69 kDa , observed, 69 kDa KDa
DDX5
DDX5; DEAD-Box Helicase 5; G17P1; HLR1;
P68; DEAD/H (Asp-Glu-Ala-Asp/His); Box
Polypeptide 5 (RNA Helicase, 68kD); DEAD
(Asp-Glu-Ala-Asp) Box; Polypeptide 5;
Probable ATP-Dependent RNA Helicase
DDX5; DEAD (Asp-Glu-Ala-Asp) Box
Helicase 5; DEAD Box Protein 5; RNA
Helicase P68; ATP-Dependent RNA Helicase
DDX5; EC 3.6.4.13; DEAD Box-5; EC 3.6.1;
HUMP68; HELR
A synthesized peptide derived from human DDX5

KD-Validated Anti-DDX5 Rabbit Monoclonal Antibody - Additional Information

Gene ID 1655 Other Names Probable ATP-dependent RNA helicase DDX5, 3.6.4.13, DEAD box protein 5, RNA helicase p68, DDX5, G17P1, HELR, HLR1

KD-Validated Anti-DDX5 Rabbit Monoclonal Antibody - Protein Information

Name DDX5

Synonyms G17P1, HELR, HLR1

Function

Involved in the alternative regulation of pre-mRNA splicing; its RNA helicase activity is necessary for increasing tau exon 10 inclusion and occurs in a RBM4-dependent manner. Binds to the tau pre-mRNA in the stem-loop region downstream of exon 10. The rate of ATP hydrolysis is highly stimulated by single-stranded RNA. Involved in transcriptional regulation; the function is independent of the RNA helicase activity. Transcriptional coactivator for androgen receptor AR but probably not ESR1. Synergizes with DDX17 and SRA1 RNA to activate MYOD1 transcriptional activity and involved in skeletal muscle differentiation. Transcriptional coactivator for p53/TP53 and involved in p53/TP53 transcriptional response to DNA damage and p53/TP53- dependent



apoptosis. Transcriptional coactivator for RUNX2 and involved in regulation of osteoblast differentiation. Acts as a transcriptional repressor in a promoter-specific manner; the function probably involves association with histone deacetylases, such as HDAC1. As component of a large PER complex is involved in the inhibition of 3' transcriptional termination of circadian target genes such as PER1 and NR1D1 and the control of the circadian rhythms.

Cellular Location

Nucleus. Nucleus, nucleolus Nucleus speckle. Cytoplasm. Note=During the G0 phase, predominantly located in the nucleus. Cytoplasmic levels increase during the G1/S phase. During the M phase, located at the vicinity of the condensed chromosomes. At G1, localizes in the cytoplasm

KD-Validated Anti-DDX5 Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

KD-Validated Anti-DDX5 Rabbit Monoclonal Antibody - Images



Western blotting analysis using anti-DDX5 antibody (Cat#AGI1197). Total cell lysates (30 µg) from various cell lines were loaded and separated by SDS-PAGE. The blot was incubated with anti-DDX5 antibody (Cat#AGI1197, 1:5,000) and HRP-conjugated goat anti-rabbit secondary antibody respectively.





Western blotting analysis using anti-DDX5 antibody (Cat#AGI1197). DDX5 expression in wild type (WT) and DDX5 shRNA knockdown (KD) HeLa cells with 30 μ g of total cell lysates. Hsp90 α serves as a loading control. The blot was incubated with anti-DDX5 antibody (Cat#AGI1197, 1:5,000) and HRP-conjugated goat anti-rabbit secondary antibody respectively.



Flow cytometric analysis of DDX5 expression in HepG2 cells using DDX5 antibody (Cat#AGI1197, 1:2,000). Green, isotype control; red, DDX5.



Immunocytochemical staining of HepG2 cells with DDX5 antibody (Cat#AGI1197, 1:1,000). Nuclei were stained blue with DAPI; DDX5 was stained magenta with Alexa Fluor® 647. Images were taken using Leica stellaris 5. Protein abundance based on laser Intensity and smart gain: High. Scale bar: 20 μ m.