

KD-Validated Anti-Phospho-eIF4E (S209) Rabbit Monoclonal Antibody
Rabbit monoclonal antibody
Catalog # AGI1376**Specification****KD-Validated Anti-Phospho-eIF4E (S209) Rabbit Monoclonal Antibody - Product Information**

Application	WB, FC, ICC
Primary Accession	P06730
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Isotype	Rabbit IgG
Calculated MW	Predicted, 25 kDa; observed, 25 kDa kDa
Gene Name	EIF4E
Aliases	EIF4E; Eukaryotic Translation Initiation Factor 4E; EIF4EL1; EIF4E1; EIF4F; MRNA Cap-Binding Protein; EIF-4F 25 kDa Subunit; EIF-4E; Eukaryotic Translation Initiation Factor 4E-Like 1; AUTS19; EIF4E; CBP
Immunogen	A synthesized peptide derived from human Phospho-eIF4E (S209)

KD-Validated Anti-Phospho-eIF4E (S209) Rabbit Monoclonal Antibody - Additional InformationGene ID **1977****Other Names**

Eukaryotic translation initiation factor 4E, eIF-4E, eIF4E, eIF-4F 25 kDa subunit, mRNA cap-binding protein, EIF4E (http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=3287 target="_blank">HGNC:3287), EIF4EL1, EIF4F

KD-Validated Anti-Phospho-eIF4E (S209) Rabbit Monoclonal Antibody - Protein Information**Name** EIF4E ([HGNC:3287](#))**Synonyms** EIF4EL1, EIF4F**Function**

Acts in the cytoplasm to initiate and regulate protein synthesis and is required in the nucleus for export of a subset of mRNAs from the nucleus to the cytoplasm which promotes processes such as RNA capping, processing and splicing (PubMed:[11606200](http://www.uniprot.org/citations/11606200), PubMed:[22578813](http://www.uniprot.org/citations/22578813), PubMed:[22684010](http://www.uniprot.org/citations/22684010), PubMed:[24335285](http://www.uniprot.org/citations/24335285), PubMed:[29987188](http://www.uniprot.org/citations/29987188)). Component

of the protein complex eIF4F, which is involved in the recognition of the mRNA cap, ATP-dependent unwinding of 5'-terminal secondary structure and recruitment of mRNA to the ribosome (By similarity). This protein recognizes and binds the 7-methylguanosine (m7G)-containing mRNA cap during an early step in the initiation of protein synthesis and facilitates ribosome binding by inducing the unwinding of the mRNAs secondary structures (PubMed:16271312, PubMed:22578813). Together with EIF4G1, antagonizes the scanning promoted by EIF1-EIF4G1 and is required for TISU translation, a process where the TISU element recognition makes scanning unnecessary (PubMed:29987188). In addition to its role in translation initiation, also acts as a regulator of translation and stability in the cytoplasm (PubMed:24335285). Component of the CYFIP1-EIF4E-FMR1 complex which binds to the mRNA cap and mediates translational repression: in the complex, EIF4E mediates the binding to the mRNA cap (By similarity). Component of a multiprotein complex that sequesters and represses translation of proneurogenic factors during neurogenesis (By similarity). In P-bodies, component of a complex that mediates the storage of translationally inactive mRNAs in the cytoplasm and prevents their degradation (PubMed:24335285). May play an important role in spermatogenesis through translational regulation of stage-specific mRNAs during germ cell development (By similarity). As well as its roles in translation, also involved in mRNA nucleocytoplasmic transport (By similarity). Its role in mRNA export from the nucleus to the cytoplasm relies on its ability to bind the m7G cap of RNAs and on the presence of the 50-nucleotide EIF4E sensitivity element (4ESE) in the 3'UTR of sensitive transcripts (By similarity). Interaction with the 4ESE is mediated by LRPPRC which binds simultaneously to both EIF4E and the 4ESE, thereby acting as a platform for assembly for the RNA export complex (By similarity). EIF4E-dependent mRNA export is independent of ongoing protein or RNA synthesis and is also NXF1-independent but is XPO1-dependent with LRPPRC interacting with XPO1 to form an EIF4E-dependent mRNA export complex (By similarity). Alters the composition of the cytoplasmic face of the nuclear pore to promote RNA export by reducing RANBP2 expression, relocalizing nucleoporin NUP214 and increasing expression of RANBP1 and RNA export factors DDX19 and GLE1 (By similarity). Promotes the nuclear export of cyclin CCND1 mRNA (By similarity). Promotes the nuclear export of NOS2/iNOS mRNA (PubMed:23471078). Promotes the nuclear export of MDM2 mRNA (PubMed:22684010). Promotes the export of additional mRNAs, including others involved in the cell cycle (By similarity). In the nucleus, binds to capped splice factor-encoding mRNAs and stimulates their nuclear export to enhance splice factor production by increasing their cytoplasmic availability to the translation machinery (By similarity). May also regulate splicing through interaction with the spliceosome in an RNA and m7G cap-dependent manner (By similarity). Also binds to some pre-mRNAs and may play a role in their recruitment to the spliceosome (By similarity). Promotes steady-state capping of a subset of coding and non-coding RNAs by mediating nuclear export of capping machinery mRNAs including RNMT, RNGTT and RAMAC to enhance their translation (By similarity). Stimulates mRNA 3'-end processing by promoting the expression of several core cleavage complex factors required for mRNA cleavage and polyadenylation, and may also have a direct effect through its interaction with the CPSF3 cleavage enzyme (By similarity). Rescues cells from apoptosis by promoting activation of serine/threonine- protein kinase AKT1 through mRNA export of NBS1 which potentiates AKT1 phosphorylation and also through mRNA export of AKT1 effectors, allowing for increased production of these proteins (By similarity).

Cellular Location

Cytoplasm, P-body. Cytoplasm. Cytoplasm, Stress granule. Nucleus. Nucleus speckle. Nucleus, nuclear body Note=Interaction with EIF4ENIF1/4E-T is required for localization to processing bodies (P-bodies) (PubMed:16157702, PubMed:24335285, PubMed:25923732). Imported in the nucleus via interaction with EIF4ENIF1/4E-T via a piggy-back mechanism (PubMed:10856257) Sequestered in the nucleus by EIF4EBP1 and EIF4EBP2 (By similarity) {ECO:0000250|UniProtKB:P63073,


ECO:0000269|PubMed:10856257, ECO:0000269|PubMed:16157702,
ECO:0000269|PubMed:24335285, ECO:0000269|PubMed:25923732 }

KD-Validated Anti-Phospho-eIF4E (S209) Rabbit Monoclonal Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


KD-Validated Anti-Phospho-eIF4E (S209) Rabbit Monoclonal Antibody - Images



Western blotting analysis using anti-phospho-eIF4E (S209) antibody (Cat#AGI1376). Total cell lysates (30 µg) from various cell lines were loaded and separated by SDS-PAGE. The blot was incubated with anti-phospho-eIF4E (S209) antibody (Cat#AGI1376, 1:10,000) and HRP-conjugated goat anti-rabbit secondary antibody respectively.




Western blotting analysis using anti-phospho-eIF4E (S209) antibody (Cat#AGI1376). Phospho-eIF4E (S209) expression in wild type (WT) and EIF4E knockdown (KD) HSHC cells with 20 µg of total cell lysates. Hsp90 α serves as a loading control. The blot was incubated with anti-phospho-eIF4E (S209) antibody (Cat#AGI1376, 1:10,000) and HRP-conjugated goat anti-rabbit secondary antibody respectively.



Copyright ©2025 Genuin Biotechnologies LLC

Phospho-eIF4E (S209)-Alexa Fluor® 647

Flow cytometric analysis of Phospho-eIF4E (S209) expression in C2C12 cells using anti-Phospho-eIF4E (S209) antibody (Cat#AGI1376, 1:2,000). Green, isotype control; red, Phospho-eIF4E (S209).



Immunocytochemical staining of C2C12 cells with Phospho-eIF4E (S209) antibody (Cat#AGI1376, 1:1,000). Nuclei were stained blue with DAPI; Phospho-eIF4E (S209) was stained magenta with Alexa Fluor® 647. Images were taken using Leica stellaris 5. Protein abundance based on laser Intensity and smart gain: High. Scale bar, 20 µm.