

KD-Validated Anti-APE1 Rabbit Monoclonal Antibody
Rabbit monoclonal antibody
Catalog # AGI1387**Specification****KD-Validated Anti-APE1 Rabbit Monoclonal Antibody - Product Information**

Application	WB, FC
Primary Accession	P27695
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Isotype	Rabbit IgG
Calculated MW	Predicted, 36 kDa , observed, 36 kDa KDa
Gene Name	APEX1
Aliases	APEX1; Apurinic/Apyrimidinic Endodeoxyribonuclease 1; REF1; HAP1; APEN; APE; APX; REF-1; APE-1; APEX; APEX Nuclease (Multifunctional DNA Repair Enzyme); DNA-(Apurinic Or Apyrimidinic Site) Endonuclease; Apurinic-Apyrimidinic Endonuclease 1; Redox Factor-1; APE1; APEX Nuclease (Multifunctional DNA Repair Enzyme); Deoxyribonuclease (Apurinic Or Apyrimidinic); Apurinic/Apyrimidinic (Antibodyasic) Endonuclease; DNA-(Apurinic Or Apyrimidinic Site) Lyase; AP Endonuclease Class I; AP Endonuclease 1; Protein REF-1; APEX Nuclease; EC 4.2.99.18; EC 3.1.11.2; AP Lyase
Immunogen	A synthesized peptide derived from human APE1

KD-Validated Anti-APE1 Rabbit Monoclonal Antibody - Additional Information

Gene ID	328
Other Names	DNA repair nuclease/redox regulator APEX1, 3.1.11.2, 3.1.21.-, APEX nuclease, APEN, Apurinic-apyrimidinic endonuclease 1, AP endonuclease 1, APE-1, DNA-(apurinic or apyrimidinic site) endonuclease, Redox factor-1, REF-1, DNA repair nuclease/redox regulator APEX1, mitochondrial, APEX1, APE, APE1, APEX, APX, HAP1, REF1

KD-Validated Anti-APE1 Rabbit Monoclonal Antibody - Protein Information**Name** APEX1**Synonyms** APE, APE1, APEX, APX, HAP1, REF1**Function**

Multifunctional protein that plays a central role in the cellular response to oxidative stress. The two

Page 2/4

href="http://www.uniprot.org/citations/18579163" target="_blank">18579163, PubMed:8355688, PubMed:9108029). Involved in calcium-dependent down-regulation of parathyroid hormone (PTH) expression by binding to negative calcium response elements (nCaREs). Together with HNRNPL or the dimer XRCC5/XRCC6, associates with nCaRE, acting as an activator of transcriptional repression (PubMed:11809897, PubMed:14633989, PubMed:8621488). May also play a role in the epigenetic regulation of gene expression by participating in DNA demethylation (PubMed:21496894). Stimulates the YBX1-mediated MDR1 promoter activity, when acetylated at Lys-6 and Lys-7, leading to drug resistance (PubMed:18809583). Plays a role in protection from granzyme-mediated cellular repair leading to cell death (PubMed:18179823). Binds DNA and RNA. Associates, together with YBX1, on the MDR1 promoter. Together with NPM1, associates with rRNA (PubMed:19188445, PubMed:19401441, PubMed:20699270).

Cellular Location


Nucleus {ECO:0000255|PROSITE-ProRule:PRU00764}. Nucleus, nucleolus. Nucleus speckle. Endoplasmic reticulum. Cytoplasm Note=Detected in the cytoplasm of B-cells stimulated to switch (By similarity). Colocalized with SIRT1 in the nucleus. Colocalized with YBX1 in nuclear speckles after genotoxic stress. Together with OGG1 is recruited to nuclear speckles in UVA-irradiated cells. Colocalized with nucleolin and NPM1 in the nucleolus. Its nucleolar localization is cell cycle dependent and requires active rRNA transcription. Colocalized with calreticulin in the endoplasmic reticulum. Translocation from the nucleus to the cytoplasm is stimulated in presence of nitric oxide (NO) and function in a CRM1-dependent manner, possibly as a consequence of demasking a nuclear export signal (amino acid position 64-80). S-nitrosylation at Cys-93 and Cys-310 regulates its nuclear-cytosolic shuttling. Ubiquitinated form is localized predominantly in the cytoplasm.


KD-Validated Anti-APE1 Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

KD-Validated Anti-APE1 Rabbit Monoclonal Antibody - Images






Western blotting analysis using anti-APE1 antibody (Cat#AGI1387). Total cell lysates (30 µg) from various cell lines were loaded and separated by SDS-PAGE. The blot was incubated with anti-APE1 antibody (Cat#AGI1387, 1:5,000) and HRP-conjugated goat anti-rabbit secondary antibody respectively.



Western blotting analysis using anti-APE1 antibody (Cat#AGI1387). APE1 expression in wild type (WT) and APE1 shRNA knockdown (KD) HeLa cells with 30 µg of total cell lysates. β-Tubulin serves as a loading control. The blot was incubated with anti-APE1 antibody (Cat#AGI1387, 1:5,000) and HRP-conjugated goat anti-rabbit secondary antibody respectively.



Flow cytometric analysis of APE1 expression in HT-1080 cells using APE1 antibody (Cat#AGI1387, 1:2,000). Green, isotype control; red, APE1.