

KD-Validated Anti-Phospho-Smad2 (S250) Rabbit Monoclonal Antibody

Rabbit monoclonal antibody Catalog # AGI1401

Specification

KD-Validated Anti-Phospho-Smad2 (S250) Rabbit Monoclonal Antibody - Product Information

Application Primary Accession Reactivity Clonality Isotype Calculated MW Gene Name Aliases	WB, FC, ICC <u>Q15796</u> Rat, Human, Mouse Monoclonal Rabbit IgG Predicted, 52 kDa , observed, 58 kDa KDa SMAD2 SMAD2; SMAD Family Member 2; JV18-1; MADR2; MADH2; Mothers Against Decapentaplegic Homolog 2; MAD Homolog 2; HMAD-2; HSMAD2; MAD, Mothers Against Decapentaplegic Homolog 2 (Drosophila); SMAD, Mothers Against DPP Homolog 2 (Drosophila); SMAD, Mothers Against DPP Homolog 2; Sma- And Mad-Related Protein 2; Mothers Against DPP Homolog 2; Mad-Related Protein 2; SMAD 2: CHTD8: Smad2: IV18: LDS6
Immunogen	A synthesized peptide derived from human Phospho-Smad2 (S250)

KD-Validated Anti-Phospho-Smad2 (S250) Rabbit Monoclonal Antibody - Additional Information

Gene ID

4087

Other Names Mothers against decapentaplegic homolog 2, MAD homolog 2, Mothers against DPP homolog 2, JV18-1, Mad-related protein 2, hMAD-2, SMAD family member 2, SMAD 2, Smad2, hSMAD2, SMAD2, MADH2, MADR2

KD-Validated Anti-Phospho-Smad2 (S250) Rabbit Monoclonal Antibody - Protein Information

Name SMAD2

Synonyms MADH2, MADR2

Function

Receptor-regulated SMAD (R-SMAD) that is an intracellular signal transducer and transcriptional modulator activated by TGF-beta (transforming growth factor) and activin type 1 receptor kinases.



Binds the TRE element in the promoter region of many genes that are regulated by TGF-beta and, on formation of the SMAD2/SMAD4 complex, activates transcription. Promotes TGFB1-mediated transcription of odontoblastic differentiation genes in dental papilla cells (By similarity). Positively regulates PDPK1 kinase activity by stimulating its dissociation from the 14-3-3 protein YWHAQ which acts as a negative regulator. May act as a tumor suppressor in colorectal carcinoma (PubMed:8752209).

Cellular Location

Cytoplasm. Nucleus. Note=Cytoplasmic and nuclear in the absence of TGF-beta. On TGF-beta stimulation, migrates to the nucleus when complexed with SMAD4 or with IPO7 (PubMed:21145499, PubMed:9865696). On dephosphorylation by phosphatase PPM1A, released from the SMAD2/SMAD4 complex, and exported out of the nucleus by interaction with RANBP1 (PubMed:16751101, PubMed:19289081). Localized mainly to the nucleus in the early stages of embryo development with expression becoming evident in the cytoplasm at the blastocyst and epiblast stages (By similarity). {ECO:0000269|PubMed:19289081, ECO:0000269|PubMed:16751101, ECO:0000269|PubMed:19289081, ECO:0000269|PubMed:21145499, ECO:0000269|PubMed:9865696}

Tissue Location

Expressed at high levels in skeletal muscle, endothelial cells, heart and placenta.

KD-Validated Anti-Phospho-Smad2 (S250) Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

KD-Validated Anti-Phospho-Smad2 (S250) Rabbit Monoclonal Antibody - Images



Western blotting analysis using anti-Phospho-Smad2 (S250) antibody (Cat#AGI1401). Total cell lysates (30 μ g) from various cell lines were loaded and separated by SDS-PAGE. The blot was incubated with anti-Phospho-Smad2 (S250) antibody (Cat#AGI1401, 1:5,000) and HRP-conjugated goat anti-rabbit secondary antibody respectively.





Western blotting analysis using anti-Phospho-Smad2 (S250) antibody (Cat#AGI1401). Phospho-Smad2 (S250) expression in wild type (WT) and Phospho-Smad2 (S250) shRNA knockdown (KD) HT-1080 cells with 30 μ g of total cell lysates. Hsp90 α serves as a loading control. The blot was incubated with anti-Phospho-Smad2 (S250) antibody (Cat#AGI1401, 1:5,000) and HRP-conjugated goat anti-rabbit secondary antibody respectively.



Flow cytometric analysis of Phospho-Smad2 (S250) expression in C2C12 cells using Phospho-Smad2 (S250) antibody (Cat#AGI1401, 1:2,000). Green, isotype control; red, Phospho-Smad2 (S250).



Immunocytochemical staining of C2C12 cells with Phospho-Smad2 (S250) antibody (Cat#AGI1401, 1:1,000). Nuclei were stained blue with DAPI; Phospho-Smad2 (S250) was stained magenta with Alexa Fluor® 647. Images were taken using Leica stellaris 5. Protein abundance based on laser Intensity and smart gain: Medium. Scale bar: 20 µm.