

#### KD-Validated Anti-Histone H3 (acetyl K56) Rabbit Monoclonal Antibody Rabbit monoclonal antibody

Catalog # AGI1439

#### Specification

# KD-Validated Anti-Histone H3 (acetyl K56) Rabbit Monoclonal Antibody - Product Information

Application Primary Accession Reactivity Clonality Isotype Calculated MW Gene Name Aliases	WB, FC, ICC <u>P68431</u> Rat, Human, Mouse Monoclonal Rabbit IgG Predicted, 15 kDa , observed, 15 kDa KDa H3C1 H3C1; H3 Clustered Histone 1; HIST1H3A; H3/A; H3FA; Histone Cluster 1 H3 Family Member A; H3 Histone Family, Member A; Histone Cluster 1, H3a; Histone 1, H3a; Histone H3.1; Histone H3/A; H3FC HIST1H3C; Histone H3/B; Histone H3/C; Histone H3/D; Histone H3/F; Histone H3/K; Histone H3/L; HIST1H3B; HIST1H3G; HIST1H3E; HIST1H3I; HIST1H3J; H3C10;
Immunogen	A synthesized peptide derived from human Histone H3 (acetyl K56)

## KD-Validated Anti-Histone H3 (acetyl K56) Rabbit Monoclonal Antibody - Additional Information

Gene ID

8350;8351;8352;8353;8354;8355;8356;835 7;8358;8968

**Other Names** Histone H3.1, Histone H3/a, Histone H3/b, Histone H3/c, Histone H3/d, Histone H3/f, Histone H3/h, Histone H3/i, Histone H3/j, Histone H3/k, Histone H3/l, H3C1 (<a href="http://www.genenames.org/cgi-bin/gene\_symbol\_report?hgnc\_id=4766" target="\_blank">HGNC:4766</a>), H3FA, HIST1H3A

## KD-Validated Anti-Histone H3 (acetyl K56) Rabbit Monoclonal Antibody - Protein Information

Name H3C1 (<u>HGNC:4766</u>)

Synonyms H3FA, HIST1H3A



Function

Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.

**Cellular Location** Nucleus. Chromosome.

### KD-Validated Anti-Histone H3 (acetyl K56) Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

#### KD-Validated Anti-Histone H3 (acetyl K56) Rabbit Monoclonal Antibody - Images



Western blotting analysis using anti-Histone H3 (acetyl K56) antibody (Cat#AGI1439). Total cell lysates (5  $\mu$ g) from various cell lines were loaded and separated by SDS-PAGE. The blot was incubated with anti-Histone H3 (acetyl K56) antibody (Cat#AGI1439, 1:5,000) and HRP-conjugated goat anti-rabbit secondary antibody respectively.

kDa 250 - 150 - 100 - 75 -	we shere	kDa 250 — 150 — 100 — 75 —	trt staat
50 <b>—</b> 37 <b>—</b>	🕳 🕳 🔶 β-Tubulin	50 <b>—</b> 37 <b>—</b>	Slotechnol
25 -		25 <b>-</b>	4 Genuin E
20 -		20 <b>—</b>	Jht @202
15 <b>—</b> 10 <b>—</b>	-	15 <b>—</b> 10 <b>—</b>	Histone H3 (acetyl K56)

Western blotting analysis using anti-Histone H3 (acetyl K56) antibody (Cat#AGI1439). Histone H3 (acetyl K56) expression in wild type (WT) and Histone H3 (acetyl K56) shRNA knockdown (KD)



HeLa cells with 5  $\mu$ g of total cell lysates.  $\beta$ -Tubulin serves as a loading control. The blot was incubated with anti-Histone H3 (acetyl K56) antibody (Cat#AGI1439, 1:5,000) and HRP-conjugated goat anti-rabbit secondary antibody respectively.



Flow cytometric analysis of Histone H3 (acetyl K56) expression in HAP-1 cells using Histone H3 (acetyl K56) antibody (Cat#AGI1439, 1:2,000). Green, isotype control; red, Histone H3 (acetyl K56).



Immunocytochemical staining of HAP-1 cells with Histone H3 (acetyl K56) antibody (Cat#AGI1439, 1:1,000). Nuclei were stained blue with DAPI; Histone H3 (acetyl K56) was stained magenta with Alexa Fluor® 647. Images were taken using Leica stellaris 5. Protein abundance based on laser Intensity and smart gain: Medium. Scale bar: 20 μm.