

KD-Validated Anti-Ku70 Rabbit Monoclonal Antibody

Rabbit monoclonal antibody
Catalog # AGI1493

Specification

KD-Validated Anti-Ku70 Rabbit Monoclonal Antibody - Product Information

Application	WB, FC, ICC
Primary Accession	P12956
Reactivity	Human
Clonality	Monoclonal
Isotype	Rabbit IgG
Calculated MW	Predicted, 70 kDa, observed, 70 kDa
Gene Name	Ku70
Aliases	XRCC6; X-Ray Repair Cross Complementing 6; X-Ray Repair Complementing Defective Repair In Chinese Hamster Cells 6; G22P1; KU70; ML8; X-Ray Repair Cross-Complementing Protein 6; Thyroid Autoantigen 70kDa (Ku Antigen); ATP-Dependent DNA Helicase 2 Subunit 1; Thyroid Autoantigen 70kD (Ku Antigen); 5'-Deoxyribose-5-Phosphate Lyase Ku70; Lupus Ku Autoantigen Protein P70; 70 kDa Subunit Of Ku Antigen; DNA Repair Protein XRCC6; Ku Autoantigen, 70kDa; 5'-DRP Lyase Ku70; D22S731; D22S671; CTC75; CTCBF; TLAA; ATP-Dependent DNA Helicase II, 70 kDa Subunit; ATP-Dependent DNA Helicase II 70 kDa Subunit; CTC Box Binding Factor 75 kDa Subunit; CTC Box-Binding Factor 75 kDa Subunit; Thyroid-Lupus Autoantigen P70; Ku Autoantigen P70 Subunit; Thyroid-Lupus Autoantigen; EC 4.2.99.-; EC 3.6.4.-; Ku70 A synthesized peptide derived from human Ku70
Immunogen	

KD-Validated Anti-Ku70 Rabbit Monoclonal Antibody - Additional Information

Gene ID	2547
Other Names	X-ray repair cross-complementing protein 6, 3.6.4.-, 4.2.99.-, 5'-deoxyribose-5-phosphate lyase Ku70, 5'-dRP lyase Ku70, 70 kDa subunit of Ku antigen, ATP-dependent DNA helicase 2 subunit 1, ATP-dependent DNA helicase II 70 kDa subunit, CTC box-binding factor 75 kDa subunit, CTC75, CTCBF, DNA repair protein XRCC6, Lupus Ku autoantigen protein p70, Ku70, Thyroid-lupus autoantigen, TLAA, X-ray repair complementing defective repair in Chinese hamster cells 6, XRCC6, G22P1

KD-Validated Anti-Ku70 Rabbit Monoclonal Antibody - Protein Information

Name XRCC6**Synonyms** G22P1**Function**

Single-stranded DNA-dependent ATP-dependent helicase that plays a key role in DNA non-homologous end joining (NHEJ) by recruiting DNA-PK to DNA (PubMed:11493912, PubMed:12145306, PubMed:20493174, PubMed:2466842, PubMed:7957065, PubMed:8621488, PubMed:9742108). Required for double-strand break repair and V(D)J recombination (PubMed:11493912, PubMed:12145306, PubMed:20493174, PubMed:2466842, PubMed:7957065, PubMed:8621488, PubMed:9742108). Also has a role in chromosome translocation (PubMed:11493912, PubMed:12145306, PubMed:20493174, PubMed:2466842, PubMed:7957065, PubMed:8621488, PubMed:9742108). Has a role in chromosome translocation (PubMed:11493912, PubMed:12145306, PubMed:20493174, PubMed:2466842, PubMed:7957065, PubMed:8621488, PubMed:9742108). The DNA helicase II complex binds preferentially to fork-like ends of double-stranded DNA in a cell cycle-dependent manner (PubMed:11493912, PubMed:12145306, PubMed:20493174, PubMed:2466842, PubMed:7957065, PubMed:8621488, PubMed:9742108). It works in the 3'-5' direction (PubMed:11493912, PubMed:12145306, PubMed:20493174, PubMed:2466842, PubMed:7957065, PubMed:8621488, PubMed:9742108). During NHEJ, the XRCC5-XRCC6 dimer performs the recognition step: it recognizes and binds to the broken ends of the DNA and protects them from further resection (PubMed:11493912, PubMed:12145306, PubMed:20493174, PubMed:2466842, PubMed:7957065, PubMed:8621488, PubMed:9742108).

<http://www.uniprot.org/citations/12145306> target="_blank">12145306, PubMed:20493174, PubMed:2466842, PubMed:7957065, PubMed:8621488, PubMed:9742108). Binding to DNA may be mediated by XRCC6 (PubMed:11493912, PubMed:12145306, PubMed:20493174, PubMed:2466842, PubMed:7957065, PubMed:8621488, PubMed:9742108). The XRCC5-XRCC6 dimer acts as a regulatory subunit of the DNA-dependent protein kinase complex DNA-PK by increasing the affinity of the catalytic subunit PRKDC to DNA by 100-fold (PubMed:11493912, PubMed:12145306, PubMed:20493174, PubMed:2466842, PubMed:7957065, PubMed:8621488, PubMed:9742108). The XRCC5-XRCC6 dimer is probably involved in stabilizing broken DNA ends and bringing them together (PubMed:11493912, PubMed:12145306, PubMed:20493174, PubMed:2466842, PubMed:7957065, PubMed:8621488, PubMed:9742108). The assembly of the DNA-PK complex to DNA ends is required for the NHEJ ligation step (PubMed:11493912, PubMed:12145306, PubMed:20493174, PubMed:2466842, PubMed:7957065, PubMed:8621488, PubMed:9742108). Probably also acts as a 5'-deoxyribose-5-phosphate lyase (5'-dRP lyase), by catalyzing the beta-elimination of the 5' deoxyribose-5-phosphate at an abasic site near double-strand breaks (PubMed:20383123). 5'-dRP lyase activity allows to 'clean' the termini of abasic sites, a class of nucleotide damage commonly associated with strand breaks, before such broken ends can be joined (PubMed:20383123). The XRCC5-XRCC6 dimer together with APEX1 acts as a negative regulator of transcription (PubMed:8621488). In association with NAA15, the XRCC5-XRCC6 dimer binds to the osteocalcin promoter and activates osteocalcin expression (PubMed:12145306). Plays a role in the regulation of DNA virus-mediated innate immune response by assembling into the HDP-RNP complex, a complex that serves as a platform for IRF3 phosphorylation and subsequent innate immune response activation through the cGAS-STING pathway (PubMed:28712728). Negatively regulates apoptosis by interacting with BAX and sequestering it from the mitochondria (PubMed:15023334). Might have deubiquitination activity, acting on BAX (PubMed:15023334).

href="http://www.uniprot.org/citations/18362350" target="_blank">18362350).

Cellular Location


Nucleus. Chromosome. Cytoplasm. Note=When trimethylated, localizes in the cytoplasm.

KD-Validated Anti-Ku70 Rabbit Monoclonal Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


KD-Validated Anti-Ku70 Rabbit Monoclonal Antibody - Images




Western blotting analysis using anti-Ku70 antibody (Cat#AGI1493). Total cell lysates (30 µg) from various cell lines were loaded and separated by SDS-PAGE. The blot was incubated with anti-Ku70 antibody (Cat#AGI1493, 1:5,000) and HRP-conjugated goat anti-rabbit secondary antibody respectively.



Western blotting analysis using anti-Ku70 antibody (Cat#AGI1493). Ku70 expression in wild type (WT) and Ku70 shRNA knockdown (KD) HeLa cells with 30 μ g of total cell lysates. β -Tubulin serves as a loading control. The blot was incubated with anti-Ku70 antibody (Cat#AGI1493, 1:5,000) and HRP-conjugated goat anti-rabbit secondary antibody respectively.



Flow cytometric analysis of Ku70 expression in HepG2 cells using Ku70 antibody (Cat#AGI1493, 1:2,000). Green, isotype control; red, Ku70.



Immunocytochemical staining of HepG2 cells with anti-Ku70 antibody (Cat #AGI1493, 1:1,000). Nuclei were stained blue with DAPI; Ku70 was stained magenta with Alexa Fluor® 647. Images were taken using Leica stellaris 5. Protein abundance based on laser Intensity and smart gain: High. Scale bar, 20 μ m.