

KD-Validated Anti-Colony Stimulating Factor 1 Rabbit Monoclonal Antibody

Rabbit monoclonal antibody Catalog # AGI1499

Specification

KD-Validated Anti-Colony Stimulating Factor 1 Rabbit Monoclonal Antibody - Product Information

Application Primary Accession Reactivity Clonality Isotype Calculated MW Gene Name Aliases	WB, FC <u>P09603</u> Human Monoclonal Rabbit IgG Predicted, 60 kDa, observed, 30 kDa KDa CSF1 CSF1; Colony Stimulating Factor 1; MCSF; M-CSF; Proteoglycan Macrophage Colony-Stimulating Factor; Colony Stimulating Factor 1 (Macrophage); Macrophage Colony Stimulating Factor 1;
Immunogen	Macrophage Colony-Stimulating Factor 1; Lanimostim; MGC31930; PG-M-CSF; CSF-1; CSF1-HERV-LTR71B Fusion Protein A synthesized peptide derived from human MCSF

KD-Validated Anti-Colony Stimulating Factor 1 Rabbit Monoclonal Antibody - Additional Information

Gene ID 1435 Other Names Macrophage colony-stimulating factor 1, CSF-1, M-CSF, MCSF, Lanimostim, Proteoglycan macrophage colony-stimulating factor, PG-M-CSF, Processed macrophage colony-stimulating factor 1, Macrophage colony-stimulating factor 1 43 kDa subunit, CSF1

KD-Validated Anti-Colony Stimulating Factor 1 Rabbit Monoclonal Antibody - Protein Information

Name CSF1

Function

Cytokine that plays an essential role in the regulation of survival, proliferation and differentiation of hematopoietic precursor cells, especially mononuclear phagocytes, such as macrophages and monocytes. Promotes the release of pro-inflammatory chemokines, and thereby plays an important role in innate immunity and in inflammatory processes. Plays an important role in the regulation of osteoclast proliferation and differentiation, the regulation of bone resorption, and is required for normal bone development. Required for normal male and female fertility. Promotes reorganization of the actin cytoskeleton, regulates formation of membrane ruffles, cell adhesion and cell migration. Plays a role in lipoprotein clearance.



Cellular Location

Cell membrane; Single-pass type I membrane protein

KD-Validated Anti-Colony Stimulating Factor 1 Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

KD-Validated Anti-Colony Stimulating Factor 1 Rabbit Monoclonal Antibody - Images



Western blotting analysis using anti-Colony stimulating factor 1 antibody (Cat#AGI1499). Total cell lysates ($30 \mu g$) from various cell lines were loaded and separated by SDS-PAGE. The blot was incubated with anti-Colony stimulating factor 1 antibody (Cat#AGI1499, 1:5,000) and HRP-conjugated goat anti-rabbit secondary antibody respectively.



Western blotting analysis using anti-Colony stimulating factor 1 antibody (Cat#AGI1499). Colony stimulating factor 1 expression in wild type (WT) and Colony stimulating factor 1 shRNA knockdown (KD) HeLa cells with 30 μ g of total cell lysates. β -Tubulin serves as a loading control. The blot was incubated with anti-Colony stimulating factor 1 antibody (Cat.#AGI1499, 1:5,000) and HRP-conjugated goat anti-rabbit secondary antibody respectively.





Flow cytometric analysis of Colony stimulating factor 1 expression in HAP-1 cells using Colony stimulating factor 1 antibody (Cat#AGI1499, 1:2,000). Green, isotype control; red, Colony stimulating factor 1.