

KD-Validated Anti-Glutamic-oxaloacetic transaminase 2 Rabbit Monoclonal Antibody

Rabbit monoclonal antibody Catalog # AGI1579

Specification

KD-Validated Anti-Glutamic-oxaloacetic transaminase 2 Rabbit Monoclonal Antibody - Product Information

Application WB, FC
Primary Accession P00505
Reactivity Rat, Human, Mouse

Clonality Monoclonal Isotype Rabbit IgG

Calculated MW Predicted, 48 kDa , observed , 43 kDa KDa

Gene Name GC

Aliases

GOT2; Glutamic-Oxaloacetic Transaminase
2; KYAT4; Kynurenine Aminotransferase IV;

MAspAT; KATIV; KAT4; Plasma

Membrane-Associated Fatty Acid-Binding

Protein; Glutamic-Oxaloacetic

Transaminase 2, Mitochondrial; Aspartate

Aminotransferase, Mitochondrial;

Kynurenine--Oxoglutarate Transaminase

IV; Kynurenine--Oxoglutarate

Transaminase 4; Glutamate Oxaloacetate

Transaminase 2; Kynurenine Aminotransferase 4; Aspartate

Aminotransferase 2; Fatty Acid-Binding Protein; Aspartate Transaminase 2; Transaminase A; EC 2.6.1.1; MitAAT; FABP-1; FABPpm; Glutamic-Oxaloacetic Transaminase 2, Mitochondrial (Aspartate Aminotransferase 2); EC 2.6.1.7; EC 2.6.1;

MITAAT; DEE82

Immunogen A synthesized peptide derived from human

GOT2

KD-Validated Anti-Glutamic-oxaloacetic transaminase 2 Rabbit Monoclonal Antibody - Additional Information

Gene ID **2806**

Other Names

Aspartate aminotransferase, mitochondrial, mAspAT, 2.6.1.1, 2.6.1.7, Fatty acid-binding protein, FABP-1, Glutamate oxaloacetate transaminase 2, Kynurenine aminotransferase 4, Kynurenine aminotransferase IV, Kynurenine--oxoglutarate transaminase 4, Kynurenine--oxoglutarate transaminase IV, Plasma membrane-associated fatty acid-binding protein, FABPpm, Transaminase A, GOT2 (HGNC:4433)

KD-Validated Anti-Glutamic-oxaloacetic transaminase 2 Rabbit Monoclonal Antibody -



Protein Information

Name GOT2 (HGNC:4433)

Function

Catalyzes the irreversible transamination of the L-tryptophan metabolite L-kynurenine to form kynurenic acid (KA). As a member of the malate-aspartate shuttle, it has a key role in the intracellular NAD(H) redox balance. Is important for metabolite exchange between mitochondria and cytosol, and for amino acid metabolism. Facilitates cellular uptake of long-chain free fatty acids.

Cellular Location


Mitochondrion matrix. Cell membrane. Note=Exposure to alcohol promotes translocation to the cell membrane.

KD-Validated Anti-Glutamic-oxaloacetic transaminase 2 Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

KD-Validated Anti-Glutamic-oxaloacetic transaminase 2 Rabbit Monoclonal Antibody - Images




Western blotting analysis using anti-Glutamic-oxaloacetic transaminase 2 antibody (Cat#AGI1579). Total cell lysates (30 μ g) from various cell lines were loaded and separated by SDS-PAGE. The blot was incubated with anti-Glutamic-oxaloacetic transaminase 2 antibody (Cat#AGI1579, 1:5,000) and HRP-conjugated goat anti-rabbit secondary antibody respectively.





Western blotting analysis using anti-Glutamic-oxaloacetic transaminase 2 antibody (Cat#AGI1579). Glutamic-oxaloacetic transaminase 2 expression in wild type (WT) and Glutamic-oxaloacetic transaminase 2 shRNA knockdown (KD) HeLa cells with 30 μg of total cell lysates. Hsp90 α serves as a loading control. The blot was incubated with anti-Glutamic-oxaloacetic transaminase 2 antibody (Cat#AGI1579, 1:5,000) and HRP-conjugated goat anti-rabbit secondary antibody respectively.



Flow cytometric analysis of Glutamic-oxaloacetic transaminase 2 expression in HepG2 cells using Glutamic-oxaloacetic transaminase 2 antibody (Cat#AGI1579, 1:2,000). Green, isotype control; red, Glutamic-oxaloacetic transaminase 2.