

KD-Validated Anti-DNA Polymerase gamma Rabbit Monoclonal Antibody
Rabbit monoclonal antibody
Catalog # AGI1650**Specification****KD-Validated Anti-DNA Polymerase gamma Rabbit Monoclonal Antibody - Product Information**

Application	WB, FC, ICC
Primary Accession	P54098
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Isotype	Rabbit IgG
Calculated MW	Predicted, 1 40 kDa , observed , 140 kDa
Gene Name	KDa
Aliases	POLG DNA Polymerase Gamma, Catalytic Subunit; POLG1; POLGA; Mitochondrial DNA Polymerase Catalytic Subunit; Polymerase (DNA) Gamma, Catalytic Subunit; Polymerase (DNA Directed), Gamma; DNA Polymerase Subunit Gamma-1; PolG-Alpha; EC 2.7.7.7; MDP1; Mitochondrial Polymerase Gamma Catalytic Subunit; MTDPS4A; MTDPS4B; MIRAS; SANDO; SCAE; PEO
Immunogen	A synthesized peptide derived from human DNA Polymerase gamma

KD-Validated Anti-DNA Polymerase gamma Rabbit Monoclonal Antibody - Additional Information

Gene ID	5428
Other Names	DNA polymerase subunit gamma-1, 2.7.7.7, 3'-5' exodeoxyribonuclease, 3.1.11.-, 5'-deoxyribose-phosphate lyase, 4.2.99.-, Mitochondrial DNA polymerase catalytic subunit, PolG-alpha, POLG {ECO:0000303 PubMed:10827171, ECO:0000312 HGNC:HGNC:9179}

KD-Validated Anti-DNA Polymerase gamma Rabbit Monoclonal Antibody - Protein Information

Name POLG {ECO:0000303|PubMed:10827171, ECO:0000312|HGNC:HGNC:9179}

Function

Catalytic subunit of DNA polymerase gamma solely responsible for replication of mitochondrial DNA (mtDNA). Replicates both heavy and light strands of the circular mtDNA genome using a single-stranded DNA template, RNA primers and the four deoxyribonucleoside triphosphates as substrates (PubMed:11477093, PubMed:<a href="http://www.uniprot.org/citations/11897778"

target="_blank">11897778, PubMed:15917273, PubMed:19837034, PubMed:9558343). Has 5' -> 3' polymerase activity. Functionally interacts with TWNK and SSBP1 at the replication fork to form a highly processive replisome, where TWNK unwinds the double-stranded DNA template prior to replication and SSBP1 covers the parental heavy strand to enable continuous replication of the entire mitochondrial genome. A single nucleotide incorporation cycle includes binding of the incoming nucleotide at the insertion site, a phosphodiester bond formation reaction that extends the 3'-end of the primer DNA, and translocation of the primer terminus to the post-insertion site. After completing replication of a mtDNA strand, mediates 3' -> 5' exonucleolytic degradation at the nick to enable proper ligation (PubMed:11477093, PubMed:11897778, PubMed:15167897, PubMed:15917273, PubMed:19837034, PubMed:26095671, PubMed:9558343). Highly accurate due to high nucleotide selectivity and 3' -> 5' exonucleolytic proofreading. Proficiently corrects base substitutions, single-base additions and deletions in non-repetitive sequences and short repeats, but displays lower proofreading activity when replicating longer homopolymeric stretches. Exerts exonuclease activity toward single-stranded DNA and double-stranded DNA containing 3'-terminal mispairs. When a misincorporation occurs, transitions from replication to a pro-nucleolytic editing mode and removes the missincorporated nucleoside in the exonuclease active site. Proceeds via an SN2 nucleolytic mechanism in which Asp-198 catalyzes phosphodiester bond hydrolysis and Glu-200 stabilizes the leaving group. As a result the primer strand becomes one nucleotide shorter and is positioned in the post-insertion site, ready to resume DNA synthesis (PubMed:10827171, PubMed:11477094, PubMed:11504725, PubMed:37202477). Exerts 5'-deoxyribose phosphate (dRP) lyase activity and mediates repair-associated mtDNA synthesis (gap filling) in base-excision repair pathway. Catalyzes the release of the 5'-terminal 2-deoxyribose-5-phosphate sugar moiety from incised apurinic/apyrimidinic (AP) sites to produce a substrate for DNA ligase. The dRP lyase reaction does not require divalent metal ions and likely proceeds via a Schiff base intermediate in a beta-elimination reaction mechanism (PubMed:9770471).

Cellular Location

Mitochondrion. Mitochondrion matrix, mitochondrion nucleoid

KD-Validated Anti-DNA Polymerase gamma Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

KD-Validated Anti-DNA Polymerase gamma Rabbit Monoclonal Antibody - Images