

**KD-Validated Anti-Ubiquitination Factor E4A Rabbit Monoclonal Antibody**  
**Rabbit monoclonal antibody**  
**Catalog # AGI1717****Specification****KD-Validated Anti-Ubiquitination Factor E4A Rabbit Monoclonal Antibody - Product Information**

Application	WB, FC, ICC
Primary Accession	<a href="#">Q14139</a>
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Isotype	Rabbit IgG
Calculated MW	Predicted, 123 kDa , observed , 123 kDa
Gene Name	UBE4A
Aliases	UBE4A; Ubiquitination Factor E4A; KIAA0126; UBOX2; UFD2; E4; Ubiquitination Factor E4A (UFD2 Homolog, Yeast); RING-Type E3 Ubiquitin Transferase E4 A; Ubiquitin Conjugation Factor E4 A; Ubiquitination Factor E4A (Homologous To Yeast UFD2); EC 2.3.2.27; NEDHMS
Immunogen	A synthesized peptide derived from human UBE4A

**KD-Validated Anti-Ubiquitination Factor E4A Rabbit Monoclonal Antibody - Additional Information**

Gene ID **9354**

**Other Names**

Ubiquitin conjugation factor E4 A, 2.3.2.27, RING-type E3 ubiquitin transferase E4 A, UBE4A (<a href="[http://www.genenames.org/cgi-bin/gene\\_symbol\\_report?hgnc\\_id=12499](http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=12499)" target="\_blank">HGNC:12499</a>)

**KD-Validated Anti-Ubiquitination Factor E4A Rabbit Monoclonal Antibody - Protein Information**

Name UBE4A ([HGNC:12499](#))

**Function**

Ubiquitin-protein ligase that probably functions as an E3 ligase in conjunction with specific E1 and E2 ligases. May also function as an E4 ligase mediating the assembly of polyubiquitin chains on substrates ubiquitinated by another E3 ubiquitin ligase. Mediates 'Lys-48'-linked polyubiquitination of substrates.

**Cellular Location**


Cytoplasm {ECO:0000250|UniProtKB:E9Q735}.

## KD-Validated Anti-Ubiquitination Factor E4A Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## KD-Validated Anti-Ubiquitination Factor E4A Rabbit Monoclonal Antibody - Images




Copyright ©2024 Genuin Biotechnology LLC

Western blotting analysis using anti-ubiquitination factor E4 A antibody (Cat#AGI1717). Total cell lysates (30 µg) from various cell lines were loaded and separated by SDS-PAGE. The blot was incubated with anti-ubiquitination factor E4 A antibody (Cat#AGI1717, 1:5,000) and HRP-conjugated goat anti-rabbit secondary antibody respectively.




Copyright ©2024 Genuin Biotechnology LLC

Western blotting analysis using anti-ubiquitination factor E4 A antibody (Cat#AGI1717). ubiquitination factor E4 A expression in wild type (WT) and ubiquitination factor E4 A shRNA knockdown (KD) HeLa cells with 20 µg of total cell lysates. β-Tubulin serves as a loading control. The blot was incubated with anti-ubiquitination factor E4 A antibody (Cat#AGI1717, 1:5,000) and HRP-conjugated goat anti-rabbit secondary antibody respectively.



Copyright ©2024 Genmab Biotechnologies LLC



Immunocytochemical staining of HepG2 cells with anti-ubiquitination factor E4 A antibody (Cat#AGI1717, 1:1,000). Nuclei were stained blue with DAPI; ubiquitination factor E4 A was stained magenta with Alexa Fluor® 647. Images were taken using Leica stellaris 5. Protein abundance based on laser Intensity and smart gain: Medium. Scale bar: 20  $\mu$ m.