

KD-Validated Anti-DSTN Mouse Monoclonal Antibody
Mouse monoclonal antibody
Catalog # AGI1907**Specification**

KD-Validated Anti-DSTN Mouse Monoclonal Antibody - Product Information

Application	WB, ICC
Primary Accession	P60981
Reactivity	Human
Clonality	Monoclonal
Isotype	Mouse IgG2a
Calculated MW	Predicted, 19 kDa, Observed, 19 kDa
Gene Name	KDa DSTN
Aliases	DSTN; Destrin, Actin Depolymerizing Factor; ACTDP; ADF; Actin-Depolymerizing Factor; Destrin; BA462D18.2 (Destrin (Actin Depolymerizing Factor ADF) (ACTDP)); Epididymis Secretory Sperm Binding Protein; Destrin (Actin Depolymerizing Factor); Epididymis Luminal Protein 32; BA462D18.2; HEL32; DSN
Immunogen	Recombinant protein of human DSTN

KD-Validated Anti-DSTN Mouse Monoclonal Antibody - Additional Information

Gene ID	11034
Other Names	
Destrin, Actin-depolymerizing factor, ADF, DSTN, ACTDP, DSN	

KD-Validated Anti-DSTN Mouse Monoclonal Antibody - Protein Information**Name** DSTN**Synonyms** ACTDP, DSN**Function**

Actin-depolymerizing protein. Severs actin filaments (F- actin) and binds to actin monomers (G-actin). Acts in a pH-independent manner.

Tissue Location


Widely distributed in various tissues.

KD-Validated Anti-DSTN Mouse Monoclonal Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


KD-Validated Anti-DSTN Mouse Monoclonal Antibody - Images



Western blotting analysis using anti-DSTN antibody (Cat#AGI1907). Total cell lysates (30 μ g) from various cell lines were loaded and separated by SDS-PAGE. The blot was incubated with anti-DSTN antibody (Cat#AGI1907, 1:2,500) and HRP-conjugated goat anti-mouse secondary antibody respectively.



Western blotting analysis using anti-DSTN antibody (Cat#AGI1907). DSTN expression in wild type (WT) and DSTN shRNA knockdown (KD) HeLa cells with 20 μ g of total cell lysates. β -Tubulin serves as a loading control. The blot was incubated with anti-DSTN antibody (Cat#AGI1907, 1:2,500) and HRP-conjugated goat anti-mouse secondary antibody respectively.



Immunocytochemical staining of HepG2 cells with anti-DSTN antibody (Cat#AGI1907, 1:1,000). Nuclei were stained blue with DAPI; DSTN was stained magenta with Alexa Fluor® 647. Images were taken using Leica stellaris 5. Protein abundance based on laser Intensity and Smart

Gain□Low. Scale bar, 20 μ m.