

KD-Validated Anti-JUN Mouse Monoclonal Antibody Mouse monoclonal antibody Catalog # AGI1916

Specification

KD-Validated Anti-JUN Mouse Monoclonal Antibody - Product Information

Application Primary Accession Reactivity Clonality Isotype Calculated MW Gene Name Aliases	WB, FC, ICC <u>P05412</u> Rat, Human, Mouse Monoclonal Mouse IgG1 Predicted, 36 kDa, Observed, 40kDa KDa JUN JUN; Jun Proto-Oncogene, AP-1 Transcription Factor Subunit; V-Jun Avian Sarcoma Virus 17 Oncogene Homolog 2; C-Jun; AP-1; Transcription Factor AP-1 Subunit Jun; Transcription Factor Jun; Proto-Oncogene C-Jun; Activator Protein 1; Jun Oncogene; AP1; P39; V-Jun Sarcoma Virus 17 Oncogene Homolog: Jun Activation
Immunogon	Virus 17 Oncogene Homolog; Jun Activation Domain Binding Protein; Enhancer-Binding Protein AP1; Transcription Factor AP-1; Proto-Oncogene CJun; CJUN Pecombinant protein of human IUN
immunogen	Recombinant protein of numan Jun

KD-Validated Anti-JUN Mouse Monoclonal Antibody - Additional Information

Gene ID3725Other NamesTranscription factor Jun, Activator protein 1, AP1, Proto-oncogene c-Jun, Transcription factor AP-1
subunit Jun, V-jun avian sarcoma virus 17 oncogene homolog, p39, JUN

KD-Validated Anti-JUN Mouse Monoclonal Antibody - Protein Information

Name JUN

Function

Transcription factor that recognizes and binds to the AP-1 consensus motif 5'-TGA[GC]TCA-3' (PubMed:10995748, PubMed:22083952). Heterodimerizes with proteins of the FOS family to form an AP-1 transcription complex, thereby enhancing its DNA binding activity to the AP-1 consensus sequence 5'-TGA[GC]TCA-3' and enhancing its transcriptional activity (By similarity). Together with FOSB, plays a role in activation-induced cell death of T cells by binding to the AP-1 promoter site of FASLG/CD95L, and inducing its transcription in response to activation of the TCR/CD3 signaling pathway (PubMed:12618758). Promotes activity of NR5A1 when phosphorylated by HIPK3 leading to increased steroidogenic gene



expression upon cAMP signaling pathway stimulation (PubMed:17210646). Involved in activated KRAS-mediated transcriptional activation of USP28 in colorectal cancer (CRC) cells (PubMed:24623306). Binds to the USP28 promoter in colorectal cancer (CRC) cells (PubMed:24623306).

Cellular Location Nucleus.

Tissue Location Expressed in the developing and adult prostate and prostate cancer cells.

KD-Validated Anti-JUN Mouse Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

KD-Validated Anti-JUN Mouse Monoclonal Antibody - Images



Western blotting analysis using anti-JUN antibody (Cat#AGI1916). Total cell lysates (30 μ g) from various cell lines were loaded and separated by SDS-PAGE. The blot was incubated with anti-JUN antibody (Cat#AGI1916, 1:2,500) and HRP-conjugated goat anti-mouse secondary antibody respectively.





Western blotting analysis using anti-JUN antibody (Cat#AGI1916). JUN expression in wild type (WT) and JUN shRNA knockdown (KD) C2C12 cells with 20 μ g of total cell lysates. Hsp90 α serves as a loading control. The blot was incubated with anti-JUN antibody (Cat#AGI1916, 1:5,000) and HRP-conjugated goat anti-mouse secondary antibody respectively.



Flow cytometric analysis of JUN expression in HT-1080 cells using anti-JUN antibody (Cat#AGI1916, 1:2,000). Green, isotype control; red, JUN.



Immunocytochemical staining of HT-1080 cells with anti-JUN antibody (Cat#AGI1916, 1:1,000). Nuclei were stained blue with DAPI; JUN was stained magenta with Alexa Fluor® 647. Images were taken using Leica stellaris 5. Protein abundance based on laser Intensity and Smart Gain[]Medium. Scale bar, 20 μ m.