

KD-Validated Anti-BID Mouse Monoclonal Antibody
Mouse monoclonal antibody
Catalog # AGI1990**Specification****KD-Validated Anti-BID Mouse Monoclonal Antibody - Product Information**

Application	WB
Primary Accession	P55957
Reactivity	Human
Clonality	Monoclonal
Isotype	Mouse IgG1 kappa
Calculated MW	Predicted, 22 kDa, observed, 19 kDa
Gene Name	KDa BID
Aliases	BID; BH3 Interacting Domain Death Agonist; BH3-Interacting Domain Death Agonist; P22 BID; Human BID Coding Sequence; Apoptic Death Agonist; Desmocollin Type 4; FP497
Immunogen	Recombinant protein of human BID

KD-Validated Anti-BID Mouse Monoclonal Antibody - Additional Information

Gene ID 637

Other Names

BH3-interacting domain death agonist, p22 BID, BID, BH3-interacting domain death agonist p15, p15 BID, BH3-interacting domain death agonist p13, p13 BID, BH3-interacting domain death agonist p11, p11 BID, BID

KD-Validated Anti-BID Mouse Monoclonal Antibody - Protein Information

Name BID

Function

Induces caspases and apoptosis (PubMed:14583606). Counters the protective effect of BCL2 (By similarity).

Cellular Location

Cytoplasm. Mitochondrion membrane. Mitochondrion outer membrane. Note=When uncleaved, it is predominantly cytoplasmic. [BH3-interacting domain death agonist p13]: Mitochondrion membrane {ECO:0000250|UniProtKB:P70444}. Note=Associated with the mitochondrial membrane. {ECO:0000250|UniProtKB:P70444} [Isoform 3]: Cytoplasm

Tissue Location


[Isoform 2]: Expressed in spleen, pancreas and placenta (at protein level). [Isoform 4]: Expressed in lung and pancreas (at protein level).

KD-Validated Anti-BID Mouse Monoclonal Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

KD-Validated Anti-BID Mouse Monoclonal Antibody - Images



Western blotting analysis using anti-BH3 interacting domain death agonist antibody (Cat#AGI1990). Total cell lysates (30 μ g) from various cell lines were loaded and separated by SDS-PAGE. The blot was incubated with anti-BH3 interacting domain death agonist antibody (Cat#AGI1990, 1:5,000) and HRP-conjugated goat anti-mouse secondary antibody respectively.



Western blotting analysis using anti-BH3 interacting domain death agonist antibody (Cat#AGI1990). BH3 interacting domain death agonist expression in wild-type (WT) and BH3 interacting domain death agonist (BID) shRNA knockdown (KD) HeLa cells with 20 μ g of total cell lysates. Hsp90 α serves as a loading control. The blot was incubated with anti-BH3 interacting domain death agonist antibody (Cat#AGI1990, 1:5,000) and HRP-conjugated goat anti-mouse secondary antibody respectively.