

KD-Validated Anti-Galectin 1 Rabbit Polyclonal Antibody
Rabbit polyclonal antibody
Catalog # AGI2126**Specification****KD-Validated Anti-Galectin 1 Rabbit Polyclonal Antibody - Product Information**

Application	WB
Primary Accession	P09382
Reactivity	Rat, Human, Mouse
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	Predicted, 15 kDa, observed, 15 kDa kDa
Gene Name	LGALS1
Aliases	LGALS1; Galectin 1; GBP; Lectin, Galactoside-Binding, Soluble, 1; Beta-Galactoside-Binding Lectin L-14-I; Putative MAPK-Activating Protein PM12; 14 kDa Laminin-Binding Protein; Lactose-Binding Lectin 1; S-Lac Lectin 1; 14 kDa Lectin; Galectin-1; Galaptin; HLBP14; Gal-1; HBL; HPL; Epididymis Secretory Sperm Binding Protein; Beta-Galactoside-Binding Protein 14kDa; Lectin Galactoside-Binding Soluble 1; GAL1 A synthesized peptide derived from human Galectin 1
Immunogen	

KD-Validated Anti-Galectin 1 Rabbit Polyclonal Antibody - Additional Information

Gene ID	3956
Other Names	
Galectin-1, Gal-1, 14 kDa laminin-binding protein, HLBP14, 14 kDa lectin, Beta-galactoside-binding lectin L-14-I, Galaptin, HBL, HPL, Lactose-binding lectin 1, Lectin galactoside-binding soluble 1, Putative MAPK-activating protein PM12, S-Lac lectin 1, LGALS1 (http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=6561)	
HGNC:6561	

KD-Validated Anti-Galectin 1 Rabbit Polyclonal Antibody - Protein Information**Name** LGALS1 ([HGNC:6561](#))**Function**

Lectin that binds beta-galactoside and a wide array of complex carbohydrates. Plays a role in regulating apoptosis, cell proliferation and cell differentiation. Inhibits CD45 protein phosphatase activity and therefore the dephosphorylation of Lyn kinase. Strong inducer of T-cell apoptosis. Plays a negative role in Th17 cell differentiation via activation of the receptor CD69 (PubMed:[24752896](http://www.uniprot.org/citations/24752896)).

Cellular Location

Secreted, extracellular space, extracellular matrix. Cytoplasm. Secreted Note=Can be secreted; the secretion is dependent on protein unfolding and facilitated by the cargo receptor TMED10; it results in protein translocation from the cytoplasm into the ERGIC (endoplasmic reticulum- Golgi intermediate compartment) followed by vesicle entry and secretion.

Tissue Location


Expressed in placenta, maternal decidua and fetal membranes. Within placenta, expressed in trophoblasts, stromal cells, villous endothelium, syncytiotrophoblast apical membrane and villous stroma. Within fetal membranes, expressed in amnion, chorioamniotic mesenchyma and chorion (at protein level). Expressed in cardiac, smooth, and skeletal muscle, neurons, thymus, kidney and hematopoietic cells.

KD-Validated Anti-Galectin 1 Rabbit Polyclonal Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

KD-Validated Anti-Galectin 1 Rabbit Polyclonal Antibody - Images



Western blotting analysis using anti-galectin 1 antibody (Cat#AGI2126). Total cell lysates (30 µg) from various cell lines were loaded and separated by SDS-PAGE. The blot was incubated with anti-galectin 1 antibody (Cat#AGI2126, 1:2,500) and HRP-conjugated goat anti-rabbit secondary antibody respectively.



Western blotting analysis using anti-galectin 1 antibody (Cat #AGI2126). Galectin 1 expression in wild-type (WT) and galectin 1 (LGALS1) shRNA knockdown (KD) HeLa cells with 20 µg of total cell lysates. Hsp90 α serves as a loading control. The blot was incubated with anti-galectin 1 antibody (Cat #AGI2126, 1:2,500) and HRP-conjugated goat anti-rabbit secondary antibody respectively.