

KD-Validated Anti-BCAM Rabbit Monoclonal Antibody
Rabbit monoclonal Antibody
Catalog # AGI2216**Specification****KD-Validated Anti-BCAM Rabbit Monoclonal Antibody - Product Information**

Application	WB
Primary Accession	P50895
Reactivity	Human
Clonality	Monoclonal
Isotype	Rabbit IgG
Calculated MW	Predicted, 67 kDa, observed, 80 kDa kDa
Gene Name	BCAM
Aliases	BCAM; Basal Cell Adhesion Molecule (Lutheran Blood Group); F8/G253; CD239; B-CAM; LU; Basal Cell Adhesion Molecule (Lu And Au Blood Groups); B-CAM Cell Surface Glycoprotein; Basal Cell Adhesion Molecule; Auberger B Antigen; F8/G253 Antigen; MSK19; Lutheran Blood Group (Auberger B Antigen Included); Lutheran Blood Group Variant LUGA; Lutheran Blood Group Glycoprotein; B-Cell Adhesion Molecule; Lutheran Antigen; CD239 Antigen; AU
Immunogen	A synthesized peptide derived from human BCAM

KD-Validated Anti-BCAM Rabbit Monoclonal Antibody - Additional Information

Gene ID 4059

Other Names

Basal cell adhesion molecule, Auberger B antigen, B-CAM cell surface glycoprotein, F8/G253 antigen, Lutheran antigen, Lutheran blood group glycoprotein, CD239, BCAM, LU, MSK19

KD-Validated Anti-BCAM Rabbit Monoclonal Antibody - Protein Information**Name** BCAM**Synonyms** LU, MSK19**Function**

Transmembrane glycoprotein that functions as both a receptor and an adhesion molecule playing a crucial role in cell adhesion, motility, migration and invasion (PubMed:9616226, PubMed:31413112). Extracellular domain enables binding to extracellular matrix proteins, such as laminin, integrin and other ligands while its intracellular domain interacts with cytoskeletal proteins like hemoglobin,

facilitating cell signal transduction (PubMed:17158232). Serves as a receptor for laminin alpha-5/LAMA5 to promote cell adhesion (PubMed:15975931). Mechanistically, JAK2 induces BCAM phosphorylation and activates its adhesion to laminin by stimulating a Rap1/AKT signaling pathway in the absence of EPOR (PubMed:23160466).

Cellular Location

Cell membrane; Single-pass type I membrane protein

Tissue Location


Wide tissue distribution (highest in the pancreas and very low in brain). Closely associated with the basal layer of cells in epithelia and the endothelium of blood vessel walls

KD-Validated Anti-BCAM Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

KD-Validated Anti-BCAM Rabbit Monoclonal Antibody - Images



Copyright ©2025 Genuin Biotechnologies LLC

Western blotting analysis using anti-CD239 antibody (Cat#65254). Total cell lysates (30 µg) from various cell lines were loaded and separated by SDS-PAGE. The blot was incubated with anti-CD239 antibody (Cat#65254, 1:5,000) and HRP-conjugated goat anti-rabbit secondary antibody (Cat#201, 1:20,000) respectively. Image was developed using NaQ™ ECL Substrate Kit (Cat#716).



Copyright ©2025 Genuin Biotechnologies LLC

Western blotting analysis using anti-CD239 antibody (Cat#65254). CD239 expression in wild-type (WT) and BCAM shRNA knockdown (KD) HeLa cells with 20 µg of total cell lysates. Hsp90 α serves as a loading control. The blot was incubated with anti-CD239 antibody (Cat#65254, 1:5,000) and HRP-conjugated goat anti-rabbit secondary antibody (Cat#201, 1:20,000) respectively. Image was developed using FeQ™ ECL Substrate Kit (Cat#226).