

KD-Validated Anti-FNTA Rabbit Monoclonal Antibody

Rabbit monoclonal antibody Catalog # AGI2237

Specification

KD-Validated Anti-FNTA Rabbit Monoclonal Antibody - Product Information

Application WB, FC Primary Accession P49354

Reactivity
Clonality
Monoclonal
Isotype
Rat, Human, Mouse
Monoclonal
Rabbit IgG

Calculated MW Predicted, 44 kDa; observed, 44 kDa KDa

Gene Name FN1

Aliases

FNTA; Farnesyltransferase, CAAX Box,
Subunit Alpha; Protein Farnesyltransferase
/Geranylgeranyltransferase Type-1 Subunit

Alpha; PGGT1A; PTAR2; FPTA; Protein Prenyltransferase Alpha Subunit Repeat

Containing 2; Ras Proteins

Prenyltransferase Subunit Alpha; Farnesyltransferase, CAAX Box, Alpha; GGTase-I-Alpha; FTase-Alpha; Type I Protein Geranyl-Geranyltransferase Alpha

Subunit; Type I Protein

Geranyl-Geranyltransferase Subunit Alpha;

Farnesyl-Protein Transferase

Alpha-Subunit; CAAX Farnesyltransferase Subunit Alpha; EC 2.5.1.58; EC 2.5.1.59 A synthesized peptide derived from human

FNŤA

KD-Validated Anti-FNTA Rabbit Monoclonal Antibody - Additional Information

Gene ID 2339

Other Names

Immunogen

Protein farnesyltransferase/geranylgeranyltransferase type-1 subunit alpha, 2.5.1.58, 2.5.1.59, CAAX farnesyltransferase subunit alpha, FTase-alpha, Ras proteins prenyltransferase subunit alpha, Type I protein geranyl-geranyltransferase subunit alpha, GGTase-I-alpha, FNTA

KD-Validated Anti-FNTA Rabbit Monoclonal Antibody - Protein Information

Name FNTA

Function

Essential subunit of both the farnesyltransferase and the geranylgeranyltransferase complex. Contributes to the transfer of a farnesyl or geranylgeranyl moiety from farnesyl or geranylgeranyl diphosphate to a cysteine at the fourth position from the C-terminus of several proteins having the C-terminal sequence Cys-aliphatic- aliphatic-X. May positively regulate neuromuscular junction




development downstream of MUSK via its function in RAC1 prenylation and activation.

KD-Validated Anti-FNTA Rabbit Monoclonal Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

KD-Validated Anti-FNTA Rabbit Monoclonal Antibody - Images




Western blotting analysis using anti-FNTA antibody (Cat#65441). Total cell lysates (30 μ g) from various cell lines were loaded and separated by SDS-PAGE. The blot was incubated with anti-FNTA antibody (Cat#65441, 1:5,000) and HRP-conjugated goat anti-rabbit secondary antibody (Cat#201, 1:20,000) respectively. Image was developed using FeQ $^{\text{m}}$ ECL Substrate Kit (Cat#226).



Western blotting analysis using anti-FNTA antibody (Cat#65441). FNTA expression in wild-type (WT) and FNTA shRNA knockdown (KD) HeLa cells with 20 μ g of total cell lysates. Hsp90 α serves as a loading control. The blot was incubated with anti-FNTA antibody (Cat#65441, 1:5,000) and



HRP-conjugated goat anti-rabbit secondary antibody (Cat#201, 1:20,000) respectively. Image was developed using NaQ $^{\text{\tiny M}}$ ECL Substrate Kit (Cat#716).



Flow cytometric analysis of AVEN expression in HepG2 cells using anti-AVEN antibody (Cat#5505, 1:2,000). Green, isotype control; red, AVEN.