

**KD-Validated Anti-KRIT1 Rabbit Monoclonal Antibody**  
**Rabbit monoclonal antibody**  
**Catalog # AGI2256****Specification****KD-Validated Anti-KRIT1 Rabbit Monoclonal Antibody - Product Information**

Application	WB
Primary Accession	<a href="#">O00522</a>
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Isotype	Rabbit IgG
Calculated MW	Predicted, 84 kDa; observed, 68-73 kDa
Gene Name	KRIT1
Aliases	KRIT1; KRIT1 Ankyrin Repeat Containing; Krev Interaction Trapped 1; CCM1; CAM; Cerebral Cavernous Malformations 1 Protein; Krev Interaction Trapped Protein 1; Ankyrin Repeat-Containing Protein Krit1; Cerebral Cavernous Malformations 1; KRIT1, Ankyrin Repeat Containing Recombinant protein of human KRIT1
Immunogen	

**KD-Validated Anti-KRIT1 Rabbit Monoclonal Antibody - Additional Information**

Gene ID	889
<b>Other Names</b>	
Krev interaction trapped protein 1, Krev interaction trapped 1, Cerebral cavernous malformations 1 protein, KRIT1, CCM1	

**KD-Validated Anti-KRIT1 Rabbit Monoclonal Antibody - Protein Information****Name** KRIT1**Synonyms** CCM1**Function**

Component of the CCM signaling pathway which is a crucial regulator of heart and vessel formation and integrity (By similarity). Negative regulator of angiogenesis. Inhibits endothelial proliferation, apoptosis, migration, lumen formation and sprouting angiogenesis in primary endothelial cells. Promotes AKT phosphorylation in a NOTCH- dependent and independent manner, and inhibits ERK1/2 phosphorylation indirectly through activation of the DELTA-NOTCH cascade. Acts in concert with CDH5 to establish and maintain correct endothelial cell polarity and vascular lumen and these effects are mediated by recruitment and activation of the Par polarity complex and RAP1B. Required for the localization of phosphorylated PRKCZ, PARD3, TIAM1 and RAP1B to the cell junction, and cell junction stabilization. Plays a role in integrin signaling via its interaction with ITGB1BP1; this prevents the interaction between ITGB1 and ITGB1BP1. Microtubule-associated protein that binds to phosphatidylinositol 4,5-bisphosphate (PIP2)-containing

membranes in a GTP-bound RAP1-dependent manner. Plays an important role in the maintenance of the intracellular reactive oxygen species (ROS) homeostasis to prevent oxidative cellular damage. Regulates the homeostasis of intracellular ROS through an antioxidant pathway involving FOXO1 and SOD2. Facilitates the down-regulation of cyclin-D1 (CCND1) levels required for cell transition from proliferative growth to quiescence by preventing the accumulation of intracellular ROS through the modulation of FOXO1 and SOD2 levels. May play a role in the regulation of macroautophagy through the down- regulation of the mTOR pathway (PubMed:<a href="http://www.uniprot.org/citations/26417067" target="\_blank">26417067</a>).

#### Cellular Location

Cytoplasm, cytoskeleton. Cell membrane; Peripheral membrane protein. Cell junction. Note=KRIT1 and CDH5 reciprocally regulate their localization to endothelial cell-cell junctions. Association with RAP1 relocates KRIT1 from microtubules to cell junction membranes. Translocates from the cytoplasm along microtubules to the cell membrane in a ITGB1BP1-dependent manner

#### Tissue Location


Low levels in brain. Very weak expression found in heart and muscle.

### KD-Validated Anti-KRIT1 Rabbit Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### KD-Validated Anti-KRIT1 Rabbit Monoclonal Antibody - Images



Copyright ©2025 Genuin Biotechnologies LLC

Western blotting analysis using anti-KRIT1 antibody (Cat#AGI2256). Total cell lysates (20 µg) from various cell lines were loaded and separated by SDS-PAGE. The blot was incubated with anti-KRIT1 antibody (Cat#AGI2256, 1:5,000) and HRP-conjugated goat anti-rabbit secondary antibody respectively.



Copyright ©2025 Genuin Biotechnologies LLC

Western blotting analysis using anti-KRIT1 antibody (Cat#AGI2256). KRIT1 expression in wild-type (WT) and KRIT1 shRNA knockdown (KD) HeLa cells with 20 µg of total cell lysates. Hsp90 α serves as a loading control. The blot was incubated with anti-KRIT1 antibody (Cat#AGI2256, 1:5,000) and HRP-conjugated goat anti-rabbit secondary antibody respectively.