

KO-Validated Anti-SRPK2 Mouse Monoclonal Antibody
Mouse monoclonal antibody
Catalog # AGI2420**Specification****KO-Validated Anti-SRPK2 Mouse Monoclonal Antibody - Product Information**

Application	WB, FC
Primary Accession	P78362
Reactivity	Rat, Human
Clonality	Monoclonal
Isotype	Mouse IgG2a
Calculated MW	Predicted, 78 kDa; observed, 125kDa KDa
Gene Name	SRPK2
Aliases	SRPK2; SRSF Protein Kinase 2; SFRS Protein Kinase 2; SFRSK2; Serine/Arginine-Rich Protein-Specific Kinase 2; Serine/Arginine-Rich Splicing Factor Kinase 2; SR-Protein-Specific Kinase 2; SR Protein Kinase 2; EC 2.7.11.1; Serine/Threonine-Protein Kinase SRPK2; Serine Kinase SRPK2
Immunogen	Recombinant protein of human SRPK2

KO-Validated Anti-SRPK2 Mouse Monoclonal Antibody - Additional Information

Gene ID	6733
Other Names	
SRSF protein kinase 2, 2.7.11.1, SFRS protein kinase 2, Serine/arginine-rich protein-specific kinase 2, SR-protein-specific kinase 2, SRSF protein kinase 2 N-terminal, SRSF protein kinase 2 C-terminal, SRPK2 {ECO:0000312 EMBL:AAH68547.1}	

KO-Validated Anti-SRPK2 Mouse Monoclonal Antibody - Protein Information**Name** SRPK2 {ECO:0000312|EMBL:AAH68547.1}**Function**

Serine/arginine-rich protein-specific kinase which specifically phosphorylates its substrates at serine residues located in regions rich in arginine/serine dipeptides, known as RS domains and is involved in the phosphorylation of SR splicing factors and the regulation of splicing (PubMed:18559500, PubMed:21056976, PubMed:9472028). Promotes neuronal apoptosis by up-regulating cyclin-D1 (CCND1) expression (PubMed:19592491). This is done by the phosphorylation of SRSF2, leading to the suppression of p53/TP53 phosphorylation thereby relieving the repressive effect of p53/TP53 on cyclin-D1 (CCND1) expression (PubMed:21205200). Phosphorylates ACIN1, and redistributes it from the nuclear speckles to the nucleoplasm, resulting

in cyclin A1 but not cyclin A2 up-regulation (PubMed:18559500). Plays an essential role in spliceosomal B complex formation via the phosphorylation of DDX23/PRP28 (PubMed:18425142). Probably by phosphorylating DDX23, leads to the suppression of incorrect R-loops formed during transcription; R-loops are composed of a DNA:RNA hybrid and the associated non-template single-stranded DNA (PubMed:28076779). Can mediate hepatitis B virus (HBV) core protein phosphorylation (PubMed:12134018). Plays a negative role in the regulation of HBV replication through a mechanism not involving the phosphorylation of the core protein but by reducing the packaging efficiency of the pregenomic RNA (pgRNA) without affecting the formation of the viral core particles (PubMed:16122776).

Cellular Location

Cytoplasm. Nucleus, nucleoplasm. Nucleus speckle. Chromosome. Note=Shuttles between the nucleus and the cytoplasm (PubMed:19592491, PubMed:21056976, PubMed:21157427) KAT5/TIP60 inhibits its nuclear translocation (PubMed:21157427) Phosphorylation at Thr-492 by PKB/AKT1 promotes nuclear translocation (PubMed:19592491). Preferentially localizes across the entire gene coding region (PubMed:28076779). During transcription, accumulates at chromatin loci where unscheduled R-loops form and colocalizes with paused 'Ser-5'-phosphorylated POLR2A/RNA polymerase II and helicase DDX23 (PubMed:28076779).

Tissue Location


Highly expressed in brain, moderately expressed in heart and skeletal muscle and at low levels in lung, liver, and kidney


KO-Validated Anti-SRPK2 Mouse Monoclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

KO-Validated Anti-SRPK2 Mouse Monoclonal Antibody - Images






Western blotting analysis using anti-SRPK2 antibody (Cat#AGI2420). Total cell lysates (30 µg) from various cell lines were loaded and separated by SDS-PAGE. The blot was incubated with anti-SRPK2 antibody (Cat#AGI2420, 1:1,000) and HRP-conjugated goat anti-mouse secondary antibody respectively.



Western blotting analysis using anti-SRPK2 antibody (Cat#AGI2420). SRPK2 expression in wild type (WT) and SRPK2 knockout (KO) HeLa cells with 30 µg of total cell lysates. Hsp90 α serves as a loading control. The blot was incubated with anti-SRPK2 antibody (Cat#AGI2420, 1:1,000) and HRP-conjugated goat anti-mouse secondary antibody respectively.



Flow cytometric analysis of SRPK2 expression in HepG2 cells using anti-SRPK2 antibody (Cat#AGI2420, 1:2,000). Green, isotype control; red, SRPK2.