

Ep-CAM / CD326 (Epithelial Marker)Mouse Monoclonal Antibody [Clone EGP40/837] Purified Mouse Monoclonal Antibody Catalog # AH10372

Specification

Ep-CAM / CD326 (Epithelial Marker)Mouse Monoclonal Antibody [Clone EGP40/837] - Product Information

Application Primary Accession Reactivity Host Clonality Isotype Calculated MW IF, FC <u>P16422</u> Human Mouse Monoclonal IgG1, kappa 40-43kDa KDa

Ep-CAM / CD326 (Epithelial Marker)Mouse Monoclonal Antibody [Clone EGP40/837] - Additional Information

Gene ID 4072

Other Names

Adenocarcinoma-associated Antigen; Cell Surface Glycoprotein Trop-1; EGP2; EGP314; EGP40; Epithelial Cell Adhesion Molecule; Epithelial Glycoprotein 314; ESA; KSA; TACD1; TACSTD1; TROP1; Tumor-associated Calcium Signal Transducer 1, ECS-1, Epidermal Surface Antigen 1, ESA1, FLOT2, Flotillin-2, M17S1 (Membrane Component, Chromosome 17, Surface Marker-1), REG-1, Reggie-1, Reggie-2

Target/Specificity Recombinant full-length human TACSTD1 protein

Application Note IF~~1:50~200<br \>FC~~1:10~50

Format 0.5ml at 100ug/ml with BSA and azide

Storage Store at 2 to 8°C.Antibody is stable for 24 months.

Precautions

Ep-CAM / CD326 (Epithelial Marker)Mouse Monoclonal Antibody [Clone EGP40/837] is for research use only and not for use in diagnostic or therapeutic procedures.

Ep-CAM / CD326 (Epithelial Marker)Mouse Monoclonal Antibody [Clone EGP40/837] - Protein Information

Name EPCAM



Synonyms GA733-2, M1S2, M4S1, MIC18, TACSTD1, TRO

Function

May act as a physical homophilic interaction molecule between intestinal epithelial cells (IECs) and intraepithelial lymphocytes (IELs) at the mucosal epithelium for providing immunological barrier as a first line of defense against mucosal infection. Plays a role in embryonic stem cells proliferation and differentiation. Up-regulates the expression of FABP5, MYC and cyclins A and E.

Cellular Location

Lateral cell membrane; Single-pass type I membrane protein. Cell junction, tight junction. Note=Colocalizes with CLDN7 at the lateral cell membrane and tight junction

Tissue Location

Highly and selectively expressed by undifferentiated rather than differentiated embryonic stem cells (ESC) Levels rapidly diminish as soon as ESC's differentiate (at protein levels). Expressed in almost all epithelial cell membranes but not on mesodermal or neural cell membranes. Found on the surface of adenocarcinoma.

Ep-CAM / CD326 (Epithelial Marker)Mouse Monoclonal Antibody [Clone EGP40/837] - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

Ep-CAM / CD326 (Epithelial Marker)Mouse Monoclonal Antibody [Clone EGP40/837] - Images



Flow Cytometric analysis of human Ep-CAM on MCF-7 Cells. Black: Cells alone; Green: Isotype Control; Red: PE-labeled Ep-CAM MAb (EGP40/837).

Ep-CAM / CD326 (Epithelial Marker)Mouse Monoclonal Antibody [Clone EGP40/837] -References

Bjork, P., Jonsson, U., Svedberg, H., Larsson, K., Lind, P., Dillner, J., Hedlund, G., Dohlsten, M. and Kalland, T. 1993. Isolation, partial characterization, and molecular cloning of a human colon adenocarcinoma cell-surface glyco- protein recognized by the C215 mouse monoclonal antibody. J.



Biol. Chem. 268: 24232-24241.