

Kappa Light Chain (B-Cell Marker) Antibody - With BSA and Azide
Mouse Monoclonal Antibody [Clone SPM558]
Catalog # AH10513**Specification****Kappa Light Chain (B-Cell Marker) Antibody - With BSA and Azide - Product Information**

Application	WB, IHC-P, IF, FC
Primary Accession	P01601
Other Accession	3514 , 449609 , P01834
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	Mouse / IgG1, kappa
Calculated MW	~22.5kDa KDa

Kappa Light Chain (B-Cell Marker) Antibody - With BSA and Azide - Additional Information**Other Names**

Ig kappa chain V-I region HK101, KV109

Application Note

WB~~1:1000
IHC-P~~N/A
IF~~1:50~200
FC~~1:10~50

Format

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage

Store at 2 to 8°C. Antibody is stable for 24 months.

Precautions

Kappa Light Chain (B-Cell Marker) Antibody - With BSA and Azide is for research use only and not for use in diagnostic or therapeutic procedures.

Kappa Light Chain (B-Cell Marker) Antibody - With BSA and Azide - Protein Information

Name IGKV1D-16 {ECO:0000303|PubMed:11549845, ECO:0000303|Ref.5}

Function

V region of the variable domain of immunoglobulin light chains that participates in the antigen recognition (PubMed: <http://www.uniprot.org/citations/24600447> target="_blank">24600447). Immunoglobulins, also known as antibodies, are membrane-bound or secreted glycoproteins produced by B lymphocytes. In the recognition phase of humoral immunity, the membrane-bound immunoglobulins serve as receptors which, upon binding of a specific antigen, trigger the clonal expansion and differentiation of B lymphocytes into

immunoglobulins- secreting plasma cells. Secreted immunoglobulins mediate the effector phase of humoral immunity, which results in the elimination of bound antigens (PubMed:20176268, PubMed:22158414). The antigen binding site is formed by the variable domain of one heavy chain, together with that of its associated light chain. Thus, each immunoglobulin has two antigen binding sites with remarkable affinity for a particular antigen. The variable domains are assembled by a process called V-(D)-J rearrangement and can then be subjected to somatic hypermutations which, after exposure to antigen and selection, allow affinity maturation for a particular antigen (PubMed:17576170, PubMed:20176268).

Cellular Location


Secreted. Cell membrane

Kappa Light Chain (B-Cell Marker) Antibody - With BSA and Azide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Kappa Light Chain (B-Cell Marker) Antibody - With BSA and Azide - Images



Formalin-fixed, paraffin-embedded human Tonsil stained with Kappa Light Chain Ab (SPM558).

Kappa Light Chain (B-Cell Marker) Antibody - With BSA and Azide - Background

This MAbs is specific to kappa light chain of immunoglobulin and shows no cross-reaction with lambda light chain or any of the five heavy chains. In mammals, the two light chains in an antibody are always identical, with only one type of light chain, kappa or lambda. The ratio of Kappa to Lambda is 70:30. However, with the occurrence of multiple myeloma or other B-cell malignancies this ratio is disturbed. Antibody to the kappa light chain is reportedly useful in the identification of leukemias, plasmacytomas, and certain non-Hodgkin's lymphomas. Demonstration of clonality in lymphoid infiltrates indicates that the infiltrate is malignant.

Kappa Light Chain (B-Cell Marker) Antibody - With BSA and Azide - References

Takahashi H et. al. Pathol Res Prac 189:300-311 (1993).2. Momose H et. al. Hum Pathol. 23:1115-1119 (1992)