

Ornithine Decarboxylase-1 (ODC-1) Antibody - With BSA and Azide
Mouse Monoclonal Antibody [Clone SPM565]
Catalog # AH10643

Specification

Ornithine Decarboxylase-1 (ODC-1) Antibody - With BSA and Azide - Product Information

Application	WB, IHC-P, IF, FC
Primary Accession	P11926
Other Accession	4953 , 467701
Reactivity	Human, Rat
Host	Mouse
Clonality	Monoclonal
Isotype	Mouse / IgG1, kappa
Calculated MW	53kDa kDa

Ornithine Decarboxylase-1 (ODC-1) Antibody - With BSA and Azide - Additional Information

Gene ID 4953

Other Names

Ornithine decarboxylase, ODC, 4.1.1.17, ODC1

Application Note

WB~1:1000
IHC-P~N/A
IF~1:50~200
FC~1:10~50

Format

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage

Store at 2 to 8°C. Antibody is stable for 24 months.

Precautions

Ornithine Decarboxylase-1 (ODC-1) Antibody - With BSA and Azide is for research use only and not for use in diagnostic or therapeutic procedures.

Ornithine Decarboxylase-1 (ODC-1) Antibody - With BSA and Azide - Protein Information

Name ODC1

Function


Catalyzes the first and rate-limiting step of polyamine biosynthesis that converts ornithine into putrescine, which is the precursor for the polyamines, spermidine and spermine. Polyamines are essential for cell proliferation and are implicated in cellular processes, ranging from DNA replication to apoptosis.

Ornithine Decarboxylase-1 (ODC-1) Antibody - With BSA and Azide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Ornithine Decarboxylase-1 (ODC-1) Antibody - With BSA and Azide - Images



Formalin-fixed, paraffin-embedded human Prostate Carcinoma stained with ODC-1 Monoclonal Antibody (SPM565)

Ornithine Decarboxylase-1 (ODC-1) Antibody - With BSA and Azide - Background

Recognizes a 53kDa protein, identified as the Ornithine Decarboxylase (ODC-1). ODC is the initial and rate-limiting enzyme in the biosynthetic pathway of polyamines and is involved in the conversion of ornithine to putrescine. The biological activity of ODC-1 is rapidly induced in response to virtually all agents known to promote cell proliferation including hormones, drugs, growth factors, mitogens, and tumor promoters. Reportedly, ODC mRNA levels are elevated in lung carcinomas as well as in colon adenomas and carcinomas. ODC activity in colorectal carcinomas is greater than those in adenomas and normal mucosa.

Ornithine Decarboxylase-1 (ODC-1) Antibody - With BSA and Azide - References

Schipper RG; Rutten RG; Sauerbeck M; Schielen WJ; Adams PJ; Kopitz J; Bohley P; Tesser GI; Verhofstad AA. Preparation and characterization of monoclonal antibodies against ornithine decarboxylase. *Journal of Immunological Methods*, 1993, 161(2):205-15. |