

ACTH (Adrenocorticotrophic Hormone) (Pituitary Marker) Antibody - With BSA and Azide
Mouse Monoclonal Antibody [Clone SPM333]
Catalog # AH10672

Specification

ACTH (Adrenocorticotrophic Hormone) (Pituitary Marker) Antibody - With BSA and Azide
- Product Information

Application	IHC-P, IF, FC
Primary Accession	P01189
Other Accession	5443 , 1897
Reactivity	Human, Mouse, Rat
Host	Mouse
Clonality	Monoclonal
Isotype	Mouse / IgG1, kappa
Calculated MW	ACTH is ~5kDa, and the POMC precursor is ~30kDa. The molecular weight of POMC depends upon isoform variation and post-translational modifications. KDa

ACTH (Adrenocorticotrophic Hormone) (Pituitary Marker) Antibody - With BSA and Azide
- Additional Information

Gene ID 5443

Other Names

Pro-opiomelanocortin, POMC, Corticotropin-lipotropin, NPP, Melanotropin gamma, Gamma-MSH, Potential peptide, Corticotropin, Adrenocorticotrophic hormone, ACTH, Melanotropin alpha, Alpha-MSH, Corticotropin-like intermediary peptide, CLIP, Lipotropin beta, Beta-LPH, Lipotropin gamma, Gamma-LPH, Melanotropin beta, Beta-MSH, Beta-endorphin, Met-enkephalin, POMC

Application Note

IHC-P~~N/A
IF~~1:50~200
FC~~1:10~50

Format

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage

Store at 2 to 8°C. Antibody is stable for 24 months.

Precautions

ACTH (Adrenocorticotrophic Hormone) (Pituitary Marker) Antibody - With BSA and Azide is for research use only and not for use in diagnostic or therapeutic procedures.

ACTH (Adrenocorticotrophic Hormone) (Pituitary Marker) Antibody - With BSA and Azide
- Protein Information

Name POMC ([HGNC:9201](#))

Function

Precursor protein of pituitary hormones that are involved in diverse physiological processes, including the regulation of energy balance, stress response, immune function and skin pigmentation. [Met-enkephalin]: Endogenous opiate.

Cellular Location

Secreted {ECO:0000250|UniProtKB:P01193}. Note=Melanocyte-stimulating hormone alpha and beta-endorphin are stored in separate granules in hypothalamic POMC neurons, suggesting that secretion may be under the control of different regulatory mechanisms {ECO:0000250|UniProtKB:P01193}

Tissue Location

ACTH and MSH are produced by the pituitary gland.

ACTH (Adrenocorticotrophic Hormone) (Pituitary Marker) Antibody - With BSA and Azide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ACTH (Adrenocorticotrophic Hormone) (Pituitary Marker) Antibody - With BSA and Azide - Images

ACTH (Adrenocorticotrophic Hormone) (Pituitary Marker) Antibody - With BSA and Azide - Background

ACTH (same as Corticotropin) is a 39 amino acid active peptide produced by the anterior pituitary. This MAb is specific to Synacthen (aa1-24 of ACTH); does not react with CLIP (aa17-39 of ACTH). POMC (pro-opiomelanocortin or corticotropin-lipotropin) is a 267 amino acid polypeptide hormone precursor that goes through extensive, tissue-specific posttranslational processing by convertases. POMC is cleaved into ten hormone chains named NPP, ACTH, alpha-MSH (Melanocyte Stimulating Hormone), beta-MSH, gamma-MSH, CLIP (corticotropin-like intermediary peptide), Lipotropin-beta, Lipotropin-gamma, beta-endorphin and Met-enkephalin. ACTH is also produced by cells of immune system (T-cells, B-cells, and macrophages) in response to stimuli associated with stress. Anti-ACTH is a useful marker in classification of pituitary tumors and the study of pituitary disease. It reacts with ACTH-producing cells (corticotrophs). It also may react with other tumors (e.g. some small cell carcinomas of the lung) causing paraneoplastic syndromes by secreting ACTH. Å

ACTH (Adrenocorticotrophic Hormone) (Pituitary Marker) Antibody - With BSA and Azide - References

Hsu DW et. al. American Journal of Pathology, 1991, 138(4):897-909