

Retinol Binding Protein-1 (RBP1) Antibody - With BSA and Azide

Mouse Monoclonal Antibody [Clone SPM442] Catalog # AH10711

Specification

Retinol Binding Protein-1 (RBP1) Antibody - With BSA and Azide - Product Information

Application ,14,3,
Primary Accession P02753

 Other Accession
 5947, 5948, 5950, 529571, P09455, P50120

 Reactivity
 Human, Mouse, Rat, Rabbit, Monkey, Goat

Host Mouse Clonality Monoclonal

Isotype Mouse / IgG1, kappa

Calculated MW 21-25kDa KDa

Retinol Binding Protein-1 (RBP1) Antibody - With BSA and Azide - Additional Information

Gene ID 5950

Other Names

Retinol-binding protein 4, Plasma retinol-binding protein, PRBP, RBP, Plasma retinol-binding protein(1-182), Plasma retinol-binding protein(1-181), Plasma retinol-binding protein(1-179), Plasma retinol-binding protein(1-176), RBP4

Format

200 μ ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage

Store at 2 to 8°C. Antibody is stable for 24 months.

Precautions

Retinol Binding Protein-1 (RBP1) Antibody - With BSA and Azide is for research use only and not for use in diagnostic or therapeutic procedures.

Retinol Binding Protein-1 (RBP1) Antibody - With BSA and Azide - Protein Information

Name RBP4

Function

Retinol-binding protein that mediates retinol transport in blood plasma (PubMed:5541771). Delivers retinol from the liver stores to the peripheral tissues (Probable). Transfers the bound all-trans retinol to STRA6, that then facilitates retinol transport across the cell membrane (PubMed:22665496).

Cellular Location

Secreted



Tissue Location


Detected in blood plasma and in urine (at protein level).

Retinol Binding Protein-1 (RBP1) Antibody - With BSA and Azide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

Retinol Binding Protein-1 (RBP1) Antibody - With BSA and Azide - Images



Formalin-fixed, paraffin-embedded human Hepatocellular Carcinoma stained with RBP1 (SPM442)

Retinol Binding Protein-1 (RBP1) Antibody - With BSA and Azide - Background

Recognizes a protein of 21kDa-25kDa, identified as retinol binding protein (RBP). Its epitope localizes between aa 74-182 of human RBP. This MAb recognizes reduced and carboxy-methylated RBP (RCM-RBP) as well as the circulatory RBP but not the native RBP, thereby suggesting that its epitope becomes accessible either on unfolding or upon binding of RBP to transthyretin (prealbumin). RBP is responsible for distributing retinol from the retinoid stores in the liver to the various target tissues. Once secreted into the blood with bound retinol, the vitamin carrier circulates complexed with transthyretin prior to vitamin delivery at the plasma membrane through a receptor-mediated mechanism.

Retinol Binding Protein-1 (RBP1) Antibody - With BSA and Azide - References

Reddy BM; Karande AA; Adiga PR. Antigenic determinants of human serum retinol binding protein as probed with monoclonal antibodies. Molecular Immunology, 1993, 30(15):1355-60