

---

**Hepatocyte Specific Antigen (Hepatocellular Marker) Antibody - With BSA and Azide  
Mouse Monoclonal Antibody [Clone SPM582 ]  
Catalog # AH10943**

**Specification**

---

**Hepatocyte Specific Antigen (Hepatocellular Marker) Antibody - With BSA and Azide -  
Product Information**

Application	,14,3,
Reactivity	Human, Dog
Host	Mouse
Clonality	Monoclonal
Isotype	Mouse / IgG1
Calculated MW	Not Known KDa

**Hepatocyte Specific Antigen (Hepatocellular Marker) Antibody - With BSA and Azide -  
Additional Information**

**Format**

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

**Storage**

Store at 2 to 8°C. Antibody is stable for 24 months.

**Precautions**

Hepatocyte Specific Antigen (Hepatocellular Marker) Antibody - With BSA and Azide is for research use only and not for use in diagnostic or therapeutic procedures.


**Hepatocyte Specific Antigen (Hepatocellular Marker) Antibody - With BSA and Azide -  
Protein Information**

**Hepatocyte Specific Antigen (Hepatocellular Marker) Antibody - With BSA and Azide -  
Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**Hepatocyte Specific Antigen (Hepatocellular Marker) Antibody - With BSA and Azide -  
Images**



Formalin-fixed, paraffin-embedded human Hepatocellular Carcinoma stained with HSA Monoclonal Antibody (SPM582).

#### **Hepatocyte Specific Antigen (Hepatocellular Marker) Antibody - With BSA and Azide - Background**

Hepatocyte Specific Antigen, also called Hepatocyte Paraffin 1 or Hep Par 1, localizes to the mitochondria of hepatocytes. It is a sensitive marker for distinguishing hepatocellular carcinomas (HCC) from other metastatic carcinomas as well as cholangio-carcinomas. HCCs occur primarily in the stomach, but they are also found in many other organs. The Hepatocyte Specific Antigen may also be a useful marker for intestinal metaplasia. Reportedly, strong expression of the Hepatocyte Specific Antigen correlates with smaller tumor size and longer patient survival. Occasionally, Hepatocyte Specific Antigen is also found in gastric carcinomas as well as in a few other non-hepatic tumors.

#### **Hepatocyte Specific Antigen (Hepatocellular Marker) Antibody - With BSA and Azide - References**

Wennerberg AE et. al. Am J Pathol 1993;143:1050-4. | Ramos-Vara, J.A., et al. Histochem 2002; J. 34: 397-401. | Fan, Z., et al. Mod. Pathol 2003; 16: 137-144, 2003