

# Fibronectin (8th Type III Repeat) Antibody - With BSA and Azide

Mouse Monoclonal Antibody [Clone 568]
Catalog # AH11226

# **Specification**

# Fibronectin (8th Type III Repeat) Antibody - With BSA and Azide - Product Information

Application IHC, IF, FC
Primary Accession P02751
Other Accession 2335, 203717
Reactivity Human
Host Mouse
Clonality Monoclonal

Isotype Mouse / IgG1, kappa

Calculated MW 220kDa (monomer); 440kDa (dimer) KDa

## Fibronectin (8th Type III Repeat) Antibody - With BSA and Azide - Additional Information

#### **Gene ID 2335**

## **Other Names**

Fibronectin, FN, Cold-insoluble globulin, CIG, Anastellin, Ugl-Y1, Ugl-Y2, Ugl-Y3, FN1, FN

# **Application Note**

<span class ="dilution\_IHC">IHC~~1:100~500</span><br \> <span class
="dilution IF">IF~~1:50~200</span><br \> <span class = "dilution FC">FC~~1:10~50</span>

#### Storage

Store at 2 to 8°C. Antibody is stable for 24 months.

#### **Precautions**

Fibronectin (8th Type III Repeat) Antibody - With BSA and Azide is for research use only and not for use in diagnostic or therapeutic procedures.

# Fibronectin (8th Type III Repeat) Antibody - With BSA and Azide - Protein Information

#### Name FN1 (<u>HGNC:3778</u>)

# Synonyms FN

### **Function**

Fibronectins bind cell surfaces and various compounds including collagen, fibrin, heparin, DNA, and actin (PubMed:<a href="http://www.uniprot.org/citations/3024962" target="\_blank">3024962</a>, PubMed:<a href="http://www.uniprot.org/citations/3593230" target="\_blank">3593230</a>, PubMed:<a href="http://www.uniprot.org/citations/3900070" target="\_blank">3900070</a>, PubMed:<a href="http://www.uniprot.org/citations/7989369" target="\_blank">7989369</a>). Fibronectins are involved in cell adhesion, cell motility, opsonization, wound healing, and maintenance of cell shape (PubMed:<a href="http://www.uniprot.org/citations/3024962" target="\_blank">3024962</a>, PubMed:<a href="http://www.uniprot.org/citations/3024962" target="\_blank">3024962</a>, PubMed:<a



href="http://www.uniprot.org/citations/3593230" target="\_blank">3593230</a>, PubMed:<a href="http://www.uniprot.org/citations/3900070" target="\_blank">3900070</a>, PubMed:<a href="http://www.uniprot.org/citations/7989369" target="\_blank">7989369</a>). Involved in osteoblast compaction through the fibronectin fibrillogenesis cell-mediated matrix assembly process, essential for osteoblast mineralization (By similarity). Participates in the regulation of type I collagen deposition by osteoblasts (By similarity). Acts as a ligand for the LILRB4 receptor, inhibiting FCGR1A/CD64-mediated monocyte activation (PubMed:<a href="http://www.uniprot.org/citations/34089617" target=" blank">34089617</a>).

### **Cellular Location**

Secreted, extracellular space, extracellular matrix. Secreted {ECO:0000250|UniProtKB:P11276}

#### **Tissue Location**


Expressed in the inner limiting membrane and around blood vessels in the retina (at protein level) (PubMed:29777959) Plasma FN (soluble dimeric form) is secreted by hepatocytes. Cellular FN (dimeric or cross-linked multimeric forms), made by fibroblasts, epithelial and other cell types, is deposited as fibrils in the extracellular matrix. Ugl-Y1, Ugl-Y2 and Ugl-Y3 are found in urine (PubMed:17614963).

## Fibronectin (8th Type III Repeat) Antibody - With BSA and Azide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- <u>Immunofluorescence</u>
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

# Fibronectin (8th Type III Repeat) Antibody - With BSA and Azide - Images



Formalin-fixed, paraffin-embedded human Renal Cell Carcinoma stained with Fibronectin Monoclonal Antibody (568).

# Fibronectin (8th Type III Repeat) Antibody - With BSA and Azide - Background

Epitope of this MAb is located in the 8th type III repeat in the cell-binding region of fibronectin. Fibronectins are disulfide-linked, dimeric glycoproteins of ~440kDa. They possess at least four binding sites for collagen, glycosaminoglycans, transglutaminase, and a cell surface receptor.





Tel: 858.875.1900 Fax: 858.875.1999

Fibronectins are involved in cell adhesion, tissue organization, and wound healing. Fibronectins are present in basement membranes, interstitial connective tissue matrix, and blood. Cellular fibronectin is widely distributed in the stroma of many malignant tumors. This MAb is excellent for staining of formalin-fixed, paraffin-embedded tissues.

# Fibronectin (8th Type III Repeat) Antibody - With BSA and Azide - References

Christensen, L., et al., APMIS, 98(7), 615-623 (1990). | Christensen, L., et al., APMIS, suppl. 26, 1-39 (1992). | Ljubimov AV et. al. Lab Investigation, 1995; 72:461-473