

TOX3 / TNRC9 Antibody - With BSA and Azide
Mouse Monoclonal Antibody [Clone TOX3/1124]
Catalog # AH11332**Specification**

TOX3 / TNRC9 Antibody - With BSA and Azide - Product Information

Application	WB, IF, FC
Primary Accession	O15405
Other Accession	27324 , 460789
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	Mouse / IgG1, kappa
Calculated MW	63kDa KDa

TOX3 / TNRC9 Antibody - With BSA and Azide - Additional Information**Gene ID** 27324**Other Names**

TOX high mobility group box family member 3, CAG trinucleotide repeat-containing gene F9 protein, Trinucleotide repeat-containing gene 9 protein, TOX3, CAGF9, TNRC9

Application Note

WB~~1:1000
IF~~1:50~200
FC~~1:10~50

Storage

Store at 2 to 8°C. Antibody is stable for 24 months.

Precautions

TOX3 / TNRC9 Antibody - With BSA and Azide is for research use only and not for use in diagnostic or therapeutic procedures.

TOX3 / TNRC9 Antibody - With BSA and Azide - Protein Information**Name** TOX3**Synonyms** CAGF9, TNRC9**Function**

Transcriptional coactivator of the p300/CBP-mediated transcription complex. Activates transactivation through cAMP response element (CRE) sites. Protects against cell death by inducing antiapoptotic and repressing pro-apoptotic transcripts. Stimulates transcription from the estrogen-responsive or BCL-2 promoters. Required for depolarization-induced transcription activation of the C-FOS promoter in neurons. Associates with chromatin to the estrogen-responsive C3 promoter region.

Cellular Location

Nucleus.


Tissue Location

Expressed mainly in epithelial cells. Expressed in the central nervous system (CNS), in the ileum and within the brain in the frontal and occipital lobe.

TOX3 / TNRC9 Antibody - With BSA and Azide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

TOX3 / TNRC9 Antibody - With BSA and Azide - Images

Western Blot Analysis(A) Recombinant protein (B) A549 (C) A431 stained with TOX3 Monoclonal Antibody (TOX3/1124).

TOX3 / TNRC9 Antibody - With BSA and Azide - Background

It recognizes a 63kDa protein, which is identified as TOX3. It contains a high mobility group (HMG)-box, which regulates Ca²⁺-dependent transcription in neurons through interaction with the cAMP-response-element-binding protein (CREB). TOX3 appears to be associated with breast cancer susceptibility and is expressed downstream of a cytoprotective cascade together with CITED1, a transcriptional regulator that does not bind directly to DNA. TOX3 is predominantly expressed in the brain and forms homodimers. TOX3 overexpression protects neuronal cells from cell death caused by endoplasmic reticulum stress or BAX overexpression through the induction of anti-apoptotic transcripts and repression of pro-apoptotic transcripts.

TOX3 / TNRC9 Antibody - With BSA and Azide - References

O'Flaherty, E., et al. 2003. TOX defines a conserved subfamily of HMG-box proteins. BMC Genomics 4: 13. | Smid, M., et al. 2006. Genes associated with breast cancer metastatic to bone. J. Clin. Oncol. 24: 2261-2267. |