

**Parathyroid Hormone (PTH) (C-Terminal) Antibody - With BSA and Azide**  
**Mouse Monoclonal Antibody [Clone PTH/1175 ]**  
**Catalog # AH12158**

**Specification**

**Parathyroid Hormone (PTH) (C-Terminal) Antibody - With BSA and Azide - Product Information**

Application	IHC, IF, FC
Primary Accession	<a href="#">P01270</a>
Other Accession	<a href="#">5741</a> , <a href="#">37045</a>
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	Mouse / IgG2b, kappa
Calculated MW	9kDa KDa

**Parathyroid Hormone (PTH) (C-Terminal) Antibody - With BSA and Azide - Additional Information**

**Gene ID** 5741

**Other Names**

Parathyroid hormone, PTH, Parathormone, Parathyrin, PTH

**Application Note**

IHC~~1:100~500  
IF~~1:50~200  
FC~~1:10~50

**Storage**

Store at 2 to 8°C. Antibody is stable for 24 months.

**Precautions**

Parathyroid Hormone (PTH) (C-Terminal) Antibody - With BSA and Azide is for research use only and not for use in diagnostic or therapeutic procedures.

**Parathyroid Hormone (PTH) (C-Terminal) Antibody - With BSA and Azide - Protein Information**

**Name** PTH {ECO:0000303|PubMed:35932760, ECO:0000312|HGNC:HGNC:9606}

**Function**

Parathyroid hormone elevates calcium level by dissolving the salts in bone and preventing their renal excretion (PubMed: [11604398](http://www.uniprot.org/citations/11604398), PubMed: [35932760](http://www.uniprot.org/citations/35932760)). Acts by binding to its receptor, PTH1R, activating G protein-coupled receptor signaling (PubMed: [18375760](http://www.uniprot.org/citations/18375760), PubMed: [35932760](http://www.uniprot.org/citations/35932760)). Stimulates [1-14C]-2-deoxy-D-glucose (2DG) transport and

glycogen synthesis in osteoblastic cells (PubMed:<a href="http://www.uniprot.org/citations/21076856" target="\_blank">21076856</a>).

#### Cellular Location


Secreted

#### Parathyroid Hormone (PTH) (C-Terminal) Antibody - With BSA and Azide - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### Parathyroid Hormone (PTH) (C-Terminal) Antibody - With BSA and Azide - Images



Formalin-fixed, paraffin-embedded human Parathyroid stained with PTH Monoclonal Antibody (PTH/1175).



Formalin-fixed, paraffin-embedded human Parathyroid stained with PTH Monoclonal Antibody (PTH/1175).

#### Parathyroid Hormone (PTH) (C-Terminal) Antibody - With BSA and Azide - Background

Epitope of this MAb maps in the C-terminus of PTH, a hormone produced by the parathyroid gland that regulates the concentration of calcium and phosphorus in extracellular fluid. This hormone elevates blood  $\text{Ca}^{2+}$  levels by dissolving the salts in bone and preventing their renal excretion. It is produced in the parathyroid gland as an 84 amino acid single chain polypeptide. It can also be secreted as N-terminal truncated fragments or C-terminal fragments after intracellular degradation, as in case of hypercalcemia. Defects in this gene are a cause of familial isolated hypoparathyroidism (FIH); also called autosomal dominant hypoparathyroidism or autosomal dominant hypocalcemia. FIH is characterized by hypocalcemia and hyperphosphatemia due to inadequate secretion of parathyroid hormone. Symptoms are seizures, tetany and cramps. FIH exist both as autosomal dominant and recessive forms of hypoparathyroidism.

#### **Parathyroid Hormone (PTH) (C-Terminal) Antibody - With BSA and Azide - References**

Watson, P.H. and Hanley, D.A. 1993. Parathyroid hormone: regulation of synthesis and secretion. Clin. Invest. Med. 16: 58-77. |