

GRP94 / HSP90B1 (Endoplasmic Reticulum Marker) Antibody - With BSA and Azide

Rat Monoclonal Antibody [Clone SPM249] Catalog # AH12456

Specification

GRP94 / HSP90B1 (Endoplasmic Reticulum Marker) Antibody - With BSA and Azide - Product Information

Application ,1,14,3,4,
Primary Accession P14625
Other Accession 7184, 192374

Reactivity Human, Mouse, Rat, Rabbit, Hamster,

Monkey, Pig, Chicken, Bovine, Sheep, Xenopus, Horse, Guinea Pig, Dog

Host Rat

Clonality Monoclonal

Isotype Rat / IgG2a, kappa

Calculated MW 94kDa KDa

GRP94 / HSP90B1 (Endoplasmic Reticulum Marker) Antibody - With BSA and Azide - Additional Information

Gene ID 7184

Other Names

Endoplasmin, 94 kDa glucose-regulated protein, GRP-94, Heat shock protein 90 kDa beta member 1, Tumor rejection antigen 1, gp96 homolog, HSP90B1, GRP94, TRA1

Storage

Store at 2 to 8°C. Antibody is stable for 24 months.

Precautions


GRP94 / HSP90B1 (Endoplasmic Reticulum Marker) Antibody - With BSA and Azide is for research use only and not for use in diagnostic or therapeutic procedures.

GRP94 / HSP90B1 (Endoplasmic Reticulum Marker) Antibody - With BSA and Azide - Protein Information

Name HSP90B1 {ECO:0000303|PubMed:39509507, ECO:0000312|HGNC:HGNC:12028}

Function

ATP-dependent chaperone involved in the processing of proteins in the endoplasmic reticulum, regulating their transport (PubMed:23572575, PubMed:39509507). Together with MESD, acts as a modulator of the Wnt pathway by promoting the folding of LRP6, a coreceptor of the canonical Wnt pathway (PubMed:23572575, PubMed:39509507). When associated with CNPY3, required for proper folding of Toll-like receptors (PubMed:<a





Tel: 858.875.1900 Fax: 858.875.1999

href="http://www.uniprot.org/citations/11584270" target=" blank">11584270). Promotes folding and trafficking of TLR4 to the cell surface (PubMed: 11584270). May participate in the unfolding of cytosolic leaderless cargos (lacking the secretion signal sequence) such as the interleukin 1/IL-1 to facilitate their translocation into the ERGIC (endoplasmic reticulum- Golgi intermediate compartment) and secretion; the translocation process is mediated by the cargo receptor TMED10 (PubMed: 32272059).

Cellular Location


Endoplasmic reticulum lumen. Sarcoplasmic reticulum lumen {ECO:0000250|UniProtKB:P41148}. Melanosome Note=Identified by mass spectrometry in melanosome fractions from stage I to stage IV.

GRP94 / HSP90B1 (Endoplasmic Reticulum Marker) Antibody - With BSA and Azide -**Protocols**

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

GRP94 / HSP90B1 (Endoplasmic Reticulum Marker) Antibody - With BSA and Azide -**Images**



Formalin-fixed, paraffin-embedded human Breast Carcinoma stained with GRP94 Monoclonal Antibody (SPM249).

GRP94 / HSP90B1 (Endoplasmic Reticulum Marker) Antibody - With BSA and Azide -Background

Recognizes a protein of 94kDa, which is identified as the glucose-regulated protein 94 (grp94) and also tumor rejection antigen (gp96). Grp94 shows a high degree of sequence homology with the heat shock protein 90 (hsp90). This MAb is highly specific to grp94 and shows minimal cross-reaction with other members of the HSP90 family. Grp s are a class of proteins unresponsive







to heat shock and are induced by glucose deprivation. Grp94 has been briefly studied as a prognostic factor in breast cancer.

GRP94 / HSP90B1 (Endoplasmic Reticulum Marker) Antibody - With BSA and Azide -References

Sorger, P.K. et al. J. Mol. Biol. 194: 341-344 (1987). | Tandon, A.K. et.al. Breast Cancer Res. and Treat. 16: 146 (1990). |