

CD2 / Lymphocyte Function Antigen 2 (LFA-2) Antibody - With BSA and Azide
Mouse Monoclonal Antibody [Clone HuLy-m1]
Catalog # AH12591

Specification

CD2 / Lymphocyte Function Antigen 2 (LFA-2) Antibody - With BSA and Azide - Product Information

Application	IF, FC
Primary Accession	P06729
Other Accession	914 , 523500
Reactivity	Human, Cat
Host	Mouse
Clonality	Monoclonal
Isotype	Mouse / IgG2b, kappa
Calculated MW	50kDa KDa

CD2 / Lymphocyte Function Antigen 2 (LFA-2) Antibody - With BSA and Azide - Additional Information

Gene ID 914

Other Names

T-cell surface antigen CD2, Erythrocyte receptor, LFA-2, LFA-3 receptor, Rosette receptor, T-cell surface antigen T11/Leu-5, CD2, CD2, SRBC

Application Note

IF~~1:50~200
FC~~1:10~50

Storage

Store at 2 to 8°C. Antibody is stable for 24 months.

Precautions

CD2 / Lymphocyte Function Antigen 2 (LFA-2) Antibody - With BSA and Azide is for research use only and not for use in diagnostic or therapeutic procedures.

CD2 / Lymphocyte Function Antigen 2 (LFA-2) Antibody - With BSA and Azide - Protein Information

Name CD2

Synonyms SRBC

Function

CD2 interacts with lymphocyte function-associated antigen CD58 (LFA-3) and CD48/BCM1 to mediate adhesion between T-cells and other cell types. CD2 is implicated in the triggering of T-cells, the cytoplasmic domain is implicated in the signaling function.

Cellular Location

Cell membrane; Single-pass type I membrane protein

Tissue Location

Expressed in natural killer cells (at protein level).

CD2 / Lymphocyte Function Antigen 2 (LFA-2) Antibody - With BSA and Azide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CD2 / Lymphocyte Function Antigen 2 (LFA-2) Antibody - With BSA and Azide - Images**CD2 / Lymphocyte Function Antigen 2 (LFA-2) Antibody - With BSA and Azide - Background**

CD2 interacts through its amino-terminal domain with the extracellular domain of CD58 (also designated CD2 ligand) to mediate cell adhesion. CD2/CD58 binding can enhance antigen-specific T cell activation. CD2 is a transmembrane glycoprotein that is expressed on peripheral blood T lymphocytes, NK cells and thymocytes. CD58 is a heavily glycosylated protein with a broad tissue distribution in hematopoietic and other cells, including endothelium. Interaction between CD2 and its counter receptor LFA3 (CD58) on opposing cells optimizes immune system recognition, thereby facilitating communication between helper T lymphocytes and antigen-presenting cells, as well as between cytolytic effectors and target cells.

CD2 / Lymphocyte Function Antigen 2 (LFA-2) Antibody - With BSA and Azide - References

Schlossman SF et al. Eds. Leukocyte Typing V, p342-352, Oxford University Press, Oxford, 1995. | Thurlow PJ, A monoclonal anti-pan-T-cell antibody. In vitro and in vivo studies. Transplantation 1983, 36(3):293-298 | Kuramochi T, et al. Characterization of feline T and B cells. Am J Vet Res 1987 Feb;48(2):183-185 | Dustin ML et al. Purified lymphocyte function-associated antigen 3 binds to CD2 and mediates T lymphocyte adhesion. J Exp Med 1987, 165:677-692 | Warren HS et al. A carbohydrate structure associated with CD15 (Lewis x) on myeloid cells is a novel ligand for human CD2. J Immunol 1996, 156(8):2866-2873. | Monostori E, et al. Activation of human T lymphocytes via the CD2 antigen results in tyrosine phosphorylation of T cell antigen receptor zeta-chains. J Immunol 1990, 144(3):1010-1014 | Lin H et al. Association of p59(fyn) with the T lymphocyte costimulatory receptor CD2. Binding of the Fyn Src homology (SH) 3 domain is regulated by the Fyn SH2 domain. J Biol Chem 1998, 273(31):19914-19921 | Marie-Cardine A et al. Recruitment of activated p56lck on endosomes of CD2-triggered T cells, colocalization with ZAP-70. J Biol Chem 1996, 271(34):20734-20739 | Rouleau M et al. CD2 induced apoptosis of peripheral T cells. Transplant Proc 1997, 29(5):2377-2378. | Bell GM and Imboden JB. CD2 and the regulation of T cell anergy. J Immunol 1995, 155(6):2805-280