

CD7 (T-Cell Leukemia Marker) Antibody - With BSA and Azide
Mouse Monoclonal Antibody [Clone HuLy-m2]
Catalog # AH12625**Specification**

CD7 (T-Cell Leukemia Marker) Antibody - With BSA and Azide - Product Information

| | |
|-------------------|--|
| Application | ,3,4, |
| Primary Accession | P09564 |
| Other Accession | 924 , 186820 |
| Reactivity | Human |
| Host | Mouse |
| Clonality | Monoclonal |
| Isotype | Mouse / IgG2a, kappa |
| Calculated MW | 40kDa kDa |

CD7 (T-Cell Leukemia Marker) Antibody - With BSA and Azide - Additional Information**Gene ID** 924**Other Names**

T-cell antigen CD7, GP40, T-cell leukemia antigen, T-cell surface antigen Leu-9, TP41, CD7, CD7

Storage

Store at 2 to 8°C. Antibody is stable for 24 months.

Precautions

CD7 (T-Cell Leukemia Marker) Antibody - With BSA and Azide is for research use only and not for use in diagnostic or therapeutic procedures.

CD7 (T-Cell Leukemia Marker) Antibody - With BSA and Azide - Protein Information**Name** CD7**Function**

Not yet known.

Cellular Location

Membrane; Single-pass type I membrane protein.

CD7 (T-Cell Leukemia Marker) Antibody - With BSA and Azide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)

- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CD7 (T-Cell Leukemia Marker) Antibody - With BSA and Azide - Images

CD7 (T-Cell Leukemia Marker) Antibody - With BSA and Azide - Background

Recognizes a protein of 40kDa, identified as CD7 (Workshop IV; Code T165). CD7 is a member of the immunoglobulin gene superfamily. Its N-terminal amino acids 1-107 are highly homologous to Ig kappa-L chains whereas the carboxyl-terminal region of the extracellular domain is proline-rich and has been postulated to form a stalk from which the Ig domain projects. CD7 is expressed on the majority of immature and mature T-lymphocytes, and T cell leukemia. It is also found on natural killer cells, a small subpopulation of normal B cells and on malignant B cells. Cross-linking surface CD7 positively modulates T cell and NK cell activity as measured by calcium fluxes, expression of adhesion molecules, cytokine secretion and proliferation. CD7 associates directly with phosphoinositol 3'-kinase. CD7 ligation induces production of D-3 phosphoinositides and tyrosine phosphorylation.

CD7 (T-Cell Leukemia Marker) Antibody - With BSA and Azide - References

Sandrin et al. In Leukocyte Typing III, McMichael AJ (ed), Oxford University Press, Oxford, 216219 (1987). | Knapp W et al. eds. Leukocyte typing IV, p341, Oxford University Press, Oxford, 1989. | Miwa H, et al biological characteristics of CD7(+) acute leukemia. Leuk Lymphoma. 1996;21:239-44. | Rabinowich H, et al. J. Immunol. 1994 153(8):3504-3513. | Emara M, et al. A human suppressor T-cell factor that inhibits T-cell replication by interaction with the IgM-Fc receptor (CD7). Hum Immunol. 1989;25(2):87-102. | Thurlow PJ, A monoclonal antibody detecting a new human T cell antigen, HuLy-m2. Transplantation 1984, 38(2):143-147. | Saxena A, et al. Biologic and clinical significance of CD7 expression in acute myeloid leukemia. Am J Hematol. 1998;58(4):278-84