

**CD7 (T-Cell Leukemia Marker) Antibody - With BSA and Azide**  
**Mouse Monoclonal Antibody [Clone 124-1D1 ]**  
**Catalog # AH12627****Specification****CD7 (T-Cell Leukemia Marker) Antibody - With BSA and Azide - Product Information**

Application	IF, FC
Primary Accession	<a href="#">P09564</a>
Other Accession	<a href="#">924</a> , <a href="#">186820</a>
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	Mouse / IgG1, kappa
Calculated MW	40kDa KDa

**CD7 (T-Cell Leukemia Marker) Antibody - With BSA and Azide - Additional Information****Gene ID** 924**Other Names**

T-cell antigen CD7, GP40, T-cell leukemia antigen, T-cell surface antigen Leu-9, TP41, CD7, CD7

**Application Note**

<span class = "dilution\_IF">IF~~1:50~200</span><br \><span class  
="dilution\_FC">FC~~1:10~50</span>

**Storage**

Store at 2 to 8°C. Antibody is stable for 24 months.

**Precautions**

CD7 (T-Cell Leukemia Marker) Antibody - With BSA and Azide is for research use only and not for use in diagnostic or therapeutic procedures.

**CD7 (T-Cell Leukemia Marker) Antibody - With BSA and Azide - Protein Information****Name** CD7**Function**

Transmembrane glycoprotein expressed by T-cells and natural killer (NK) cells and their precursors (PubMed:<a href="http://www.uniprot.org/citations/7506726" target="\_blank">7506726</a>). Plays a costimulatory role in T-cell activation upon binding to its ligand K12/SECTM1 (PubMed:<a href="http://www.uniprot.org/citations/10652336" target="\_blank">10652336</a>). In turn, mediates the production of cytokines such as IL-2 (PubMed:<a href="http://www.uniprot.org/citations/1709867" target="\_blank">1709867</a>). On resting NK-cells, CD7 activation results in a significant induction of interferon-gamma levels (PubMed:<a href="http://www.uniprot.org/citations/7506726" target="\_blank">7506726</a>).

**Cellular Location**

Membrane; Single-pass type I membrane protein.

**Tissue Location**

Expressed on T-cells and natural killer (NK) cells and their precursors.

**CD7 (T-Cell Leukemia Marker) Antibody - With BSA and Azide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**CD7 (T-Cell Leukemia Marker) Antibody - With BSA and Azide - Images****CD7 (T-Cell Leukemia Marker) Antibody - With BSA and Azide - Background**

Recognizes a protein of 40kDa, identified as CD7 (Workshop IV; Code T155). CD7 is a member of the immunoglobulin gene superfamily. Its N-terminal amino acids 1-107 are highly homologous to Ig kappa-L chains whereas the carboxyl-terminal region of the extracellular domain is proline-rich and has been postulated to form a stalk from which the Ig domain projects. CD7 is expressed on the majority of immature and mature T-lymphocytes, and T cell leukemia. It is also found on natural killer cells, a small subpopulation of normal B cells and on malignant B cells. Cross-linking surface CD7 positively modulates T cell and NK cell activity as measured by calcium fluxes, expression of adhesion molecules, cytokine secretion and proliferation. CD7 associates directly with phosphoinositol 3'-kinase. CD7 ligation induces production of D-3 phosphoinositides and tyrosine phosphorylation.

**CD7 (T-Cell Leukemia Marker) Antibody - With BSA and Azide - References**

Knapp W et al. eds. Leukocyte typing IV, p341, Oxford University Press, Oxford, 1989 | Miwa H, et al. Biological characteristics of CD7(+) acute leukemia. Leuk. Lymphoma. 1996, 21(3-4):239-244. Rabinowich H, et al. Signaling via CD7 molecules on human NK cells. Induction of tyrosine phosphorylation and beta 1 integrin-mediated adhesion to fibronectin J. Immunol. 1994;153(8):3504-3513. | Saxena A, et al. Biologic and clinical significance of CD7 expression in acute myeloid leukemia. Am J Hematol. 1998, 58(4):278-84